

Vol. I.

No. 1.

THE  
NATIONAL GEOGRAPHIC  
MAGAZINE.



ESTABLISHED IN 1888

NATIONAL GEOGRAPHIC SOCIETY.

WASHINGTON, D. C.

1890

Price 50 cents

## NOTICE

The River and Valleys of North Wales: William Morris Davis . . . . .	Page 145
(Illustrated by two maps and twenty-five cuts.)	
Topographic Symbols: Charles D. Walcott . . . . .	171
(Illustrated by ten plates.)	
Statistical Geography: England—Continued of Abington . . . . .	200
International Literary Contest to be held at Madrid, Spain . . . . .	221
1906, 1907	

THE  
NATIONAL GEOGRAPHIC MAGAZINE.

---

Vol. I.

1889.

No. 3.

---

THE RIVERS AND VALLEYS OF PENNSYLVANIA.\*

BY WILLIAM MORRIS DAVIS.

"In Faltensystemen von sehr hohem Alter wurde die ursprüngliche Anordnung der Langenthäler durch das Ueberhandnehmen der transversalen Erosionsfurchen oft ganz und gar verwischt."

LÖWL. Petermann's Mittheilungen, xxviii, 1883, 411.

CONTENTS.

PART FIRST. *Introductory.*

1. Plan of work here proposed.
2. General description of the topography of Pennsylvania.
3. The drainage of Pennsylvania.
4. Previous studies of Appalachian drainage.

PART SECOND. *Outline of the geological history of the region.*

5. Conditions of formation.
6. Former extension of strata to the southeast.
7. Cambro-Silurian and Permian deformations.
8. Perm-Triassic denudation.
9. Newark deposition.
10. Jurassic tilting.
11. Jura-Cretaceous denudation.
12. Tertiary elevation and denudation.
13. Later changes of level.
14. Illustrations of Pennsylvanian topography.

\* The substance of this essay was presented to the Society in a lecture on February 9th, 1889, but since then it has been much expanded.

**PART THIRD.** *General conception of the history of a river.*

15. The complete cycle of river life: youth, adolescence, maturity and old age.
16. Mutual adjustment of river courses.
17. Terminology of rivers changed by adjustment.
18. Examples of adjustments.
19. Revival of rivers by elevation and drowning by depression.
20. Opportunity for new adjustments with revival.
21. Antecedent and superimposed rivers.
22. Simple, compound, composite and complex rivers.

**PART FOURTH.** *The development of the rivers of Pennsylvania.*

23. Means of distinguishing between antecedent and adjusted consequent rivers.
24. Postulates of the argument.
25. Constructional Permian topography and consequent drainage.
26. The Jura mountains homologous with the Permian Alleghanias.
27. Development and adjustment of the Permian drainage.
28. Lateral water-gaps near the apex of synclinal ridges.
29. Departure of the Juniata from the Juniata-Catawissa syncline.
30. Avoidance of the Broad Top basin by the Juniata headwaters.
31. Reversal of larger rivers to southeast courses.
32. Capture of the Anthracite headwaters by the growing Susquehanna.
33. Present outward drainage of the Anthracite basins.
34. Homologies of the Susquehanna and Juniata.
35. Superimposition of the Susquehanna on two synclinal ridges.
36. Evidence of superimposition in the Susquehanna tributaries.
37. Events of the Tertiary cycle.
38. Tertiary adjustment of the Juniata on the Medina anticlines.
39. Migration of the Atlantic-Ohio divide.
40. Other examples of adjustments.
41. Events of the Quaternary cycle.
42. Doubtful cases.
43. Complicated history of our actual rivers.
44. Provisional conclusions.

**PART FIRST.** *Introductory.*

1. *Plan of work here proposed.*—No one now regards a river and its valley as ready-made features of the earth's surface. All are convinced that rivers have come to be what they are by slow processes of natural development, in which every peculiarity of river-course and valley-form has its appropriate cause. Being

fully persuaded of the gradual and systematic evolution of topographic forms, it is now desired, in studying the rivers and valleys of Pennsylvania, to seek the causes of the location of the streams in their present courses; to go back if possible to the early date when central Pennsylvania was first raised above the sea and trace the development of the several river systems then implanted upon it from their ancient beginning to the present time.

The existing topography and drainage system of the State will first be briefly described. We must next inquire into the geological structure of the region, follow at least in a general way the deformations and changes of attitude and altitude that it has suffered, and consider the amount of denudation that has been accomplished on its surface. We must at the same time bear in mind the natural history of rivers, their morphology and development; we must recognize the varying activities of a river in its youth and old age, the adjustments of its adolescence and maturity, and the revival of its decrepit powers when the land that it drains is elevated and it enters a new cycle of life. Finally we shall attempt to follow out the development of the rivers of Pennsylvania by applying the general principles of river history to the special case of Pennsylvania structure.

2. *General description of the topography of Pennsylvania.*—The strongly marked topographic districts of Pennsylvania can hardly be better described than by quoting the account given over a century ago by Lewis Evans, of Philadelphia, in his "Analysis of a map of the middle British colonies in America" (1755), which is as valuable from its appreciative perception as it is interesting from its early date. The following paragraphs are selected from his early pages:

"The land southwestward of Hudson's River is more regularly divided and into a greater number of stages than the other. The first object worthy of regard in this part is a rief or vein of rocks of the talky or singlassy kind, some two or three or half a dozen miles broad; rising generally some small matter higher than the adjoining land; and extending from New York city southwesterly by the lower falls of Delaware, Schuylkill, Susquehanna, Gun-Powder, Patapsco, Potomack, Rapahannock, James river and Roanoke. This was the ancient maritime boundary of America and forms a very regular curve. The land between this rief and the sea and from the Navesink hills southwest . . . may be denominated the Lower Plains, and consists of soil washed down from above and sand accumulated from the ocean. Where

these plains are not penetrated by rivers, they are a white sea-sand, about twenty feet deep and perfectly barren, as no mixture of soil helps to enrich them. But the borders of the rivers, which descend from the uplands, are rendered fertile by the soil washed down with the floods and mixt with the sands gathered from the sea. The substratum of sea-mud, shells and other foreign subjects is a perfect confirmation of this supposition. And hence it is that for 40 or 50 miles inland and all the way from the Navesink to Cape Florida, all is a perfect barren where the wash from the uplands has not enriched the borders of the rivers; or some ponds and defiles have not furnished proper support for the growth of white cedar. . . .

"From this reef of rocks, over which all the rivers fall, to that chain of broken hills, called the South mountain, there is the distance of 50, 60 or 70 miles of very uneven ground, rising sensibly as you advance further inland, and may be denominated the Upland. This consists of veins of different kinds of soil and substrata some scores of miles in length; and in some places overlaid with little ridges and chains of hills. The declivity of the whole gives great rapidity to the streams; and our violent gusts of rain have washed it all into gullies, and carried down the soil to enrich the borders of the rivers in the Lower Plains. These inequalities render half the country not easily capable of culture, and impoverishes it, where torn up by the plow, by daily washing away the richer mould that covers the surface.

"The South mountain is not in ridges like the Endless mountains, but in small, broken, steep, stoney hills; nor does it run with so much regularity. In some places it gradually degenerates to nothing, not to appear again for some miles, and in others it spreads several miles in breadth. Between South mountain and the hither chain of the Endless mountains (often for distinction called the North mountain, and in some places the Kittatinny and Pequén), there is a valley of pretty even good land, some 8, 10 or 20 miles wide, and is the most considerable quantity of valuable land that the English are possess of; and runs through New Jersey, Pennsylvania, Maryland and Virginia. It has yet obtained no general name, but may properly enough be called *Piement*, from its situation. Besides conveniences always attending good land, this valley is everywhere enriched with Limestone.

"The Endless mountains, so called from a translation of the Indian name bearing that signification, come next in order. They are not confusedly scattered and in lofty peaks overtopping one another, but stretch in long uniform ridges scarce half a mile perpendicular in any place above the intermediate valleys. Their name is expressive of their extent, though no doubt not in a literal sense. . . . The mountains are almost all so many ridges with even tops and nearly of a height. To look from these hills into the lower lands is but, as it were, into an ocean of woods, swelled and deprest here and there by little inequalities, not to be distinguished one part from another any more than the



waves of the real ocean. The uniformity of these mountains, though depriving us of an advantage in this respect, makes some amends in another. They are very regular in their courses, and confine the creeks and rivers that run between; and if we know where the gaps are that let through these streams, we are not at a loss to lay down their most considerable inflections, . . . . .

"To the northwestward of the Endless mountains is a country of vast extent, and in a manner as high as the mountains themselves. To look at the abrupt termination of it, near the sea level, as is the case on the west side of Hudson's river below Albany, it looks as a vast high mountain; for the Kaons Killa, though of more lofty stature than any other mountains in these parts of America, are but the continuation of the Plains on the top, and the cliffs of them in the front they present towards Kinderhook. These Upper Plains are of extraordinary rich level land, and extend from the Mohocks river through the country of the Confederates.\* Their termination northward is at a little distance from Lake Ontario; but what it is westward is not known, for those most extensive plains of Ohio are part of them."

These several districts recognized by Evans may be summarized as the coastal plain, of nearly horizontal Cretaceous and later beds, just entering the southeastern corner of Pennsylvania; the marginal upland of contactd schists of disputed age; the South Mountain belt of ancient and much disturbed crystalline rocks, commonly called Archean; a space between these two traversed by the sandstone lowland of the Newark formation of the great Appalachian valley of crowded Cambrian limestones and slates; the region of the even-crested, linear Paleozoic ridges, bounded by Kittatinny or Blue mountain on the southeast and by Alleghany mountain on the northwest, this being the area with which we are here most concerned; and finally the Alleghany plateau, consisting of nearly horizontal Devonian and Carboniferous beds and embracing all the western part of the state. The whole region presents the most emphatic expression not only of its structure but also of the more recent cycles of development through which it has passed. Fig. 1 represents the stronger ridges and larger streams of the greater part of the central district: it is reproduced from the expressive Topographic Map of Pennsylvania (1871) by Lesley. The Susquehanna flows down the middle, receiving the West Branch from Lock Haven

\* Referring to the league of Indian tribes, so-called.

† Russell has lately recommended the revival of this term, proposed many years ago by Redfield, as a non-committal name for the "New red sandstones" of our Atlantic slope, commonly called Triassic.




FIG. 1. Part of Topographic Map of Pontus, by J. P. Loubet (1911).



and Williamsport, the East Branch from Wilkes-Barre in the Wyoming basin, and the Juniata from the Broad Top region, south of Huntingdon. The Anthracite basins lie on the right, enclosed by zigzag ridges of Pocono and Pottsville sandstone; the Plateau, trenched by the West Branch of the Susquehanna is in the northwest. Medina sandstone forms most of the central ridges.

3. *The drainage of Pennsylvania.*—The greater part of the Alleghany plateau is drained westward into the Ohio, and with this we shall have little to do. The remainder of the plateau drainage reaches the Atlantic by two rivers, the Delaware and the Susquehanna, of which the latter is the more special object of our study. The North and West Branches of the Susquehanna rise in the plateau, which they traverse in deep valleys; thence they enter the district of the central ranges, where they unite and flow in broad lowlands among the even-crested ridges. The Juniata brings the drainage of the Broad Top region to the main stream just before their confluent current cuts across the marginal Blue Mountain. The rock-rimmed basins of the anthracite region are drained by small branches of the Susquehanna northward and westward, and by the Schuylkill and Lehigh to the south and east. The Delaware, which traverses the plateau between the Anthracite region and the Catskill Mountain front, together with the Lehigh, the Schuylkill, the little Swatara and the Susquehanna, cut the Blue Mountain by fine water-gaps, and cross the great limestone valley. The Lehigh then turns eastward and joins the Delaware, and the Swatara turns westward to the Susquehanna; but the Delaware, Schuylkill and Susquehanna all continue across South Mountain and the Newark belt, and into the low plateaus of schists beyond. The Schuylkill unites with the Delaware near Philadelphia, just below the inner margin of the coastal plain; the Delaware and the Susquehanna continue in their deflected estuaries to the sea. All of these rivers and many of their side streams are at present sunk in small valleys of moderate depth and width, below the general surface of the lowlands, and are more or less complicated with terrace gravels.

4. *Previous studies of Appalachian drainage.*—There have been no special studies of the history of the rivers of Pennsylvania in the light of what is now known of river development. A few recent essays of rather general character as far as our rivers are concerned, may be mentioned.

Peschel examined our rivers chiefly by means of general maps with little regard to the structure and complicated history of the region. He concluded that the several transverse rivers which break through the mountains, namely, the Delaware, Susquehanna and Potomac, are guided by fractures, anterior to the origin of the rivers.\* There does not seem to be sufficient evidence to support this obsolescent view, for most of the water-gaps are located independently of fractures; nor can Peschel's method of river study be trusted as leading to safe conclusions.

Tietze regards our transverse valleys as antecedent;† but this was made only as a general suggestion, for his examination of the structure and development of the region is too brief to establish this and exclude other views.‡

Löwl questions the conclusion reached by Tietze and ascribes the transverse gaps to the backward or headwater erosion of external streams, a process which he has done much to bring into its present important position, and which for him replaces the persistence of antecedent streams of other authors.§

A brief article§ that I wrote in comment on Löwl's first essay several years ago now seems to me insufficient in its method. It exaggerated the importance of antecedent streams; it took no sufficient account of the several cycles of erosion through which the region has certainly passed; and it neglected due consideration of the readjustment of initial immature stream courses during more advanced river-life. Since then, a few words in Löwl's essay have come to have more and more significance to me; he says that in mountain systems of very great age, the original arrangement of the longitudinal valleys often becomes entirely confused by means of their conquest by transverse erosion gaps. This suggestion has been so profitable to me that I have placed the original sentence at the beginning of this paper. Its thesis is the essential element of my present study.

Phillipson refers to the above-mentioned authors and gives a brief account of the arrangement of drainage areas within our Appalachians, but briefly dismisses the subject.¶ His essay contains a serviceable bibliography.

If these several earlier essays have not reached any precise

\* *Physische Erdkunde*, 1880, II, 412.

† *Jahrbuch Geol. Reichsanstalt*, XVIII, 1878, 306.

‡ *Pet. Mitth.*, 1883, 405; *Ueber Thalbildung*, Prag, 1884.

§ *Origin of Cross-valleys*. *Science*, I, 1883, 225.

¶ *Studien über Wasserschelden*. Leipzig, 1888, 146.

conclusion, it may perhaps be because the details of the geological structure and development of Pennsylvania have not been sufficiently examined. Indeed, unless the reader has already become familiar with the geological maps and reports of the Pennsylvania surveys and is somewhat acquainted with its geography, I shall hardly hope to make my case clear to him. The volumes that should be most carefully studied are, first, the always inspiring *Essay on the Geology and its Topography* (1854), by Lesley in which the immediate relation of our topography to the underlying structure is so finely described; the Geological Map of Pennsylvania (1858), the result of the labors of the first survey of the state; and the Geological Atlas of Counties, Volume X of the second survey (1885). Besides these, the numerous volumes of the final report of the first survey and numerous reports or separate counties by the second survey should be examined, as they contain many accounts of the topography although saying very little about its development. If, in addition to all this, the reader has seen the central district of the state and marvelled at its even-crested, straight and zigzag ridges, and walked through its narrow water-gaps into the enclosed coasts that Lory drama, he may then still better follow the considerations here presented.

*PART SECOND. Outline of the geological history of the region.*

§ *Conditions of formation.* The region in which the Susqueanna and the neighboring rivers are now located is built in chief part of marine sediments derived in paleozoic time from a large and now to the southwest, whose northwest coast-line probably crossed Pennsylvania somewhere in the southeastern part of the state, doubtless varying its position, however, by many miles as the sea advanced and receded in accordance with the changes in the relative altitudes of the land and water surfaces, such as have been discussed by Newberry and Chappin. The sediments thus accumulated are of enormous thickness, measuring twenty or thirty thousand feet from their crystalline foundation to the uppermost layer now remaining. The whole mass is essentially conformable in the central part of the state. Some of the formations are resistant, and these have determined the position of our ridges; others are weaker and are chosen as the sites of valleys and low lands. The first are the Onondaga and Medina sandstones, which will be here generally referred to under the latter name alone, the Pottsville conglomerate, the Juniata sandstone, the Allegheny shale, and the Westmoreland shale, the latter two of which

the whole series of bedded formations was deposited, and the basal sandstone that is generally associated with it. Whenever we now see these harder rocks, they rise above the surrounding lowland surface. On the other hand, the weaker beds are the Cambrian limestones (Trenton) and slates (Hudson River), all the Silurian except the Medina above named, the whole of the Devonian—in which however there are two hard beds of subordinate value—the Oriskany sandstone and a Chemung sandstone and conglomerate, that form low and broken ridges over the softer ground on either side of them—and the Carboniferous (Mauch Chunk red shales and some of the weaker sandstones (Coal measures).

6. *Former extension of strata to the southeast.*—We are not much concerned with the conditions under which this great series of beds was formed; but, as we appear now, it is important for us to recognize that the present southeastern margin of the beds is not by any means the original margin in that direction. It is probable that the whole mass of deposits, with greater or less variations of thickness, extended at least twenty miles beyond that of Blue Mountain, and that many of the beds extended much farther. The reason for this conclusion is a simple one. The several resistant beds above-mentioned consist of quartz sand and pebbles that cannot be derived from the underlying beds of limestones and shales; their only known source lay in the crystalline rocks of the paleozoic land to the southeast. South Mountain may possibly have made part of this paleozoic land; but it seems more probable that it was land only during the earlier Archean age, and that it was submerged and buried in Cambrian time and not again brought to the light of day until it had been crushed into many local anticlines\* whose crests were uncovered by Permian and later erosion. The occurrence of Cambrian limestone on the other side of South Mountain, taken with its complicated anticlinal structure, makes it likely that Medina time found this crystalline area entirely covered by the Cambrian beds. Medina sands must therefore have come from farther still to the southeast. A similar argument applies to the source of the Pottsville and Pottsville beds. The measure of twenty miles as the former southeastern extension of the paleozoic formations therefore seems to be a moderate one for the average of the whole series—perhaps forty would be nearer the truth.

\* See also as below.

7. *Cambro-Silurian and Permian deformations.*—This great series of once horizontal beds is now wonderfully distorted; but the distortions follow a general rule of trending northeast and southwest, and of diminishing in intensity from southeast to northwest. In the Hudson Valley, it is well known that a considerable disturbance occurred between Cambrian and Silurian time, for there the Medon lies unconformably on the Hudson River shales. It seems likely, for reasons that will be briefly given later on, that the same disturbance extended into Pennsylvania and farther southwest, but that it affected only the southeastern corner of the State; and that the unconformities in evidence of it, which are preserved in the Hudson Valley, are here lost by subsequent erosion. Wastes of the ancient land and a Cambro-Silurian annex still continued and furnished vast beds of sandstone and sandy shales to the remaining marine area, until at last the subsiding Paleozoic basin was filled up and the coal marshes extended broadly across it. At this time we may picture the drainage of the southeastern land area wandering rather slowly across the great Carboniferous plain to the still submerged basin far to the west, a condition of things that is not imperfectly represented, although in a somewhat more advanced stage, by the existing drainage of the mountains of the Appalachians across the more modern coastal plain to the Atlantic.

This condition was interrupted by the great Permian deformation that gave rise to the main ranges of the Appalachians in Pennsylvania, Virginia and Tennessee. The Permian name seems appropriate here, for while the deformation may have begun at an earlier date, and may have continued into Triassic time, its culmination seems to have been within Permian limits. It was characterized by a relentless force of compression, exerted in a southeast-northwest line, in obedience to which the whole series of the Carboniferous beds, even twenty or more thousand feet in thickness, was crowded gradually into great and small folds, trending northeast and southwest. The subjacent Archean terrane doubtless shared more or less in the disturbance. For example, South Mountain is described by Leakey as "not one mountain, but a series of mountains separated by valleys. It is, properly speaking, a system of anticlines with troughs between."<sup>\*</sup>

It appears that the South Mountain range ends eastward [in Chester and York Counties] in a part with five [artificial] ranges."

<sup>\*</sup> Proc. Amer. Acad. Sci., xii., 1878, 6.



It may be concluded with fair probability that the folds began to rise in the southeast, where they are crowded closest together, some of them having begun here while coal marshes were still forming farther west; and that the last folds to be begun were the fainter ones on the plateau, now seen in Negro mountain and Chestnut and Laurel ridges. In consequence of the inequalities in the force of compression or in the resistance of the yielding mass, the folds do not continue indefinitely with horizontal axes, but vary in height, rising or falling away in great variety. Several adjacent folds often follow some general control in this respect, four axes rising and falling together. It is to an unequal folding of this kind that we owe the location of the Anthracite ~~system in the Pennsylvania and West Virginia~~ now worn away from the prolongation of the synclines, when rise in the direction.

8. *Perm-Triassic denudation.*—During and for a long time after this period of mountain growth, the destructive processes of erosion wasted the land and covered its surface. An enormous amount of material was thus swept away and laid down in some unknown ocean bed. We shall speak of this as the Perm-Triassic period of erosion. A measure of its vast accomplishment is seen when we find that the Newark formation, which is generally correlated with Triassic or Jurassic time, lies unconformably on the eroded surface of Cambrian and Archean rocks in the southeastern part of the State, where we have concluded that the Paleozoic series once existed; where the strata must have risen at a great mountain time as a result of the Appalachian deformations; and where they must therefore have been denuded before the deposition of the Newark beds. Not only so; the moderate smoothness of the southeastern or outer boundary of the Newark formation indicates clearly enough that the surface on which that portion of the formation lies is one of so great relief or inequality; and such a surface can be carved out of an elevated land only after long continued denudation, by which topographic development is carried beyond the time of its greatest strength or maturity into the fainter expression of old age. This is a matter of some importance in our study of the development of the rivers of Pennsylvania; and it also constitutes a good part of the evidence already referred to as indicating that there must have been some earlier deformations of importance in the southeastern part of the State, for it is hardly conceivable that the great Paleozoic

mass could have been so deeply worn off of the Newark belt between the making of the last of the coal beds and the first of the Newark. It seems more in accordance with the facts here presented and with the teachings of geological history in general to suppose, as we have seen, that something of the present deformation of the ancient rocks underlying the Newark beds was given at an early date, such as that of the Green Mountain growth; and that a certain amount of the removal of the folded beds was thus made possible in another Paleozoic time; then again at some later date, as Permian, a second period of mountain growth arrived, and further folding was effected and after this came deeper erosion, thus dividing the destructive work that was done into several parts, instead of crowding it all into the post-Carboniferous time or partly assigned to it. It is indeed not impossible that an important share of what we have called the Permian deformation was, as above suggested, accomplished in the southeastern part of the State where the coal beds were yet forming at the west, many grades of sand in the sandstones of the Coal Measures may have had several temporary harts in other sandstone beds between the time of their first erosion from the Archean rocks and the much later time when they found the resting place in the Newark group.\*

3. *Newark deposition.* After the great Paleozoic and Permian-Triassic erosions thus indicated, when the southeastern area of ancient mountains had been well worn down and the Permian rocks of the central district had acquired a well developed drainage, there appeared an opportunity for great deposition in the new depression of a north-south-west belt of the deeply wasted area, across the southeastern part of the State, and into this trough the deposition, the waste from the adjacent areas on either side was carried, building the Newark formation. This may be referred to as the Newark or Trias-Jurassic period of deposition. The volume of this formation is unknown, as its thickness and original area are still undetermined, but it is pretty surely of many thousand feet in vertical measure, and its original area may have been one-fifth or a quarter in extent of its present area, if not larger yet. So great a local accumulation seems to indicate that while the best of deposition was

\* These considerations may have value in showing that the time in which the actual crushing of the Appalachians was accomplished was not so brief as is stated by Beede in a recent article in the *American Geologist* 31, 1909, 106.

sinking, the adjacent areas were rising, in order to furnish a continual supply of material, the occurrence of heavy conglomerates among the strata of the Newark basin is a strong indication of deposition, and the heavy breccias near Reading indicate the occurrence of a strong topography and a strong transporting agent to the northwest of this part of the Newark belt. It will be necessary, when the development of the ancestors of our present rivers is taken up, to consider the effects of the depression that determined the locus of Newark deposition and of the adjacent elevation that maintained a supply of material.

11. *Jurassic tilting*—Newark deposition was stopped by a gradual reversal of the conditions that introduced it. The depression of the Newark belt was after a time reversed into elevation, and the material that was being deposited was swept of the region was carried away to some unknown resting place. This disturbance, which may be regarded as a revival of the Permian activity, continued in Jurassic, or at least in post-Newark time, and resulted in the production of the angular unconformal attitude of the formation; and as far as I can correlate with the topographic map of the region, it involved there an over pushing of the curved folds of the Archaean and Paleozoic rocks. This is illustrated in figs 2 and 3,



FIG. 2.

FIG. 3.

in which the original and disturbed attitudes of the Newark and the underlying formations are roughly shown, the over-arching of the fundamental folds causing the unconformal and probably faulted structure in the overlying beds.\* If this be true, we might suspect that the asymmetrical attitude of the Appalachian folds, noted by Rogers as a characteristic of the range, is a feature that was introduced if not originated in Jurassic and not in Permian time.

\* *Amer. Journ. Science*, XXXI, 1856, 823; and *Seventh Ann. Rept. U. S. Geol. Survey*, 1884, 403.

It is not to be supposed that the Jurassic deformation was limited to the area of the Newark beds; it may have extended some way on either side; but it presumably faded out at no great distance, for it has not been noticed in the history of the Atlantic and Mississippi regions remote from the Newark belt. In the district of the central folds of Pennsylvania, with which we are particularly concerned, this deformation was probably expressed in a further folding and overpushing of the already partly folded beds, with rapidly decreasing effect to the north-west; and perhaps also by *el p-faults*, which at the surface of the *crust* and *sediments* and *hypothetical* to a high degree. The essential point for our subsequent consideration is that the Jurassic deformation was probably accompanied by a moderate elevation, for it allowed the erosion of the Newark beds and of laterally adjacent areas as well.

11. *Jura-Cretaceous denudation*.—In consequence of this elevation, a new cycle of erosion was entered upon, which I shall call the *Jura-Cretaceous cycle*. It allowed the accomplishment of

of denudation, a wide area of first relief, whose elevated remnants are now to be seen in the even ridge-crests that so strongly characterize the central district, as well as in certain other even uplands, now etched by the erosion of a later cycle of destructive work. I shall not here take space for the elaborate statement of the argument leading to this end, but its elements are as follows: the extraordinary persistent concordance among the crest-line *trilobes* of many *Medusa* and *Calymene* ridges in the central district, the generally corresponding elevation of the western plateau surface, itself a surface of erosion, and now treacheried by relatively deep and narrow valleys, the generally uniform and consistent altitude of the upland in the crystalline highlands of northern New Jersey and in the South Mountains of Pennsylvania, and the extension of the same general surface, descending slowly eastward, over the even crest-lines of the Newark trap ridges. Besides the evidence of recent central elevation thus deduced from the topography, it may be noted that a lower stand of the land in Cretaceous time than now is indicated by the erosion that the Cretaceous beds have suffered as consequence of the elevation that followed their deposition. The Cretaceous transgression in the western states doubtless opens on the problem

also. Finally it may be fairly urged that it is more accordant with what is known about old mountains in general, to suppose that their mass has stood at different attitudes with respect to base level during their long period of denudation than to suppose that they have held one attitude through all the time since their formation.

It is natural enough that the former maintenance of some lower attitude than the present should have expression in the form of the country, if it is now distinguished by subsequent erosion. It is simply the reverse of this statement that leads us to the above-stated conclusion. We may be sure that the long maintained period of relative quiet was of great importance in allowing time for the mature adjustment of the rivers of the region, and hence due account must be taken of it in a later section. I say relative quiet, for there were certainly subordinate oscillations of greater or less value. Mr. Lee has detected records of one of these about the beginning of Cretaceous time, but its effects are not now known to be of geographic value; that is, they do not now manifest themselves in the form of the present surface of the land, not only in the manner of deposition and the extinction of certain deposits\*. Another subordinate oscillation in the sense of a moderate depression seems to have extended through middle and later Cretaceous time, resulting in an in and transgression of the sea and the deposit of the Cretaceous formation unconformably on the previous land surface for a considerable distance beyond the present margin of the formation†. This is important as affecting our rivers. Although these oscillations were of considerable geological value, I do not think that, for the present purposes, they call for any primary division of the assumed tectonic cycle, for as the result of this longer period of denudation we find but a single record in the great lowland of erosion above described, a record of prime importance in the geographic development of our region, that will often be referred to. The surface of fairly relief then occupied may be called the Cretaceous surface, lowland. It may be pictured as a low, undulating plain of wide extent, with a portion of its Atlantic margin submerged and covered over with a relatively thin marine deposit of sands, clays and coals.

\* *Amer. Jour. Science*, 22: 27-38, 397-417.

† This statement is based on a study of the geographic evolution of northern New Jersey in perspective for publication.



2. *Tertiary elevation and denudation.*—This broad lowland is a lowland no longer. It has been raised over the greater part of its area into a highland, with an elevation of from one to three thousand feet, sloping gently seaward and descending under the Atlantic level near the present margin of the Cretaceous formation. The elevation seems to have taken place early in Tertiary time and will be referred to as of that date. Opportunity was then given for the revival of the previously exhausted forces of denudation, and as a consequence we now see the formerly even surface of the plain greatly roughened by the incision of deep valleys and the opening of broad lowlands on the softer rocks. Only the harder rocks remain to indicate one of the even surfaces which once stretched continuously across the whole area. The best indication of the average altitude at which the mass stood through the greater part of post-Cretaceous time is to be found on the weak shales of the Newark formation in New Jersey and Pennsylvania, and on the weak Cambrian limestones of the great Kalamazoo valley; for both of these areas have been actually again *hammered* again in the Tertiary cycle. They will be referred to as the Tertiary baselvel lowlands; and the valleys corresponding to them, cut in the harder rocks, as well as the rolling lowlands between the ridges of the central district of Pennsylvania will be regarded as of the same date. Whatever variations of level occurred in this cycle of development do not seem to have left marks of importance on the inland surface, though they may have had greater significance near the coast.

13. *Later changes of level.*—Again at the close of Tertiary time, there was an elevation of moderate amount, and to this may be referred the trenches that are so distinctly cut across the Tertiary baselvel lowland by the larger rivers, as well as the lateral



channel channels of the smaller streams. This will be called the Quaternary cycle; and for the present no farther mention of the oscillations known to have occurred in this division of time need be considered; the reader may find careful discussion of



FIG. 4.

tion in the paper by Metree, above referred to. It is proper that I should add that the suggestion of base-leveling both of the crest-lands and of the lowlands, that I have found so profitable in this and other work, is due largely to personal conference with Messrs. Holden and Metree of the Geological Survey; but it is not desired to make them in any way responsible for the statements here given.

14. *Illustrations of Pennsylvania topography*.—A few sketches made during a recent recess-trip with several students through Pennsylvania may be introduced in this connection. The first, fig. 4, is a view from Jenny Jump mountain, on the north-western side of the New Jersey highlands, looking northwest across the Katantany valley-lowland to Blue or Kittatany mountain, where it is cut at the Delaware Water-gap. The extraordinary over-crest of the mountain preserves record of the Cretaceous base-level lowland, since the elevation of the present lowland, its softer rocks have, as it were, been etched out, leaving the harder ones in relief; thus the present valley-lowland is to be explained. In consequence of the still later elevation of less amount, the Delaware has cut a trench in the present lowland, which is partly seen to the left in the sketch. Fig. 5 is a general view of the Lehigh plateau and cañon, looking south from Bald Mountain just above Penn Haven Junction. Blue mountain is the most distant crest, seen for a little space. The ridges near and above Mauch



vated, the up and surface must have been an even plain—the limestone has been lowland elevated into a plateau. The valleys

cut into the plateau during the Tertiary cycle are narrow here because the rocks are mostly hard. The steep slopes of the cañon-like valley of the Lehugh and the even crests of the ridges manifestly belong to different cycles of development. Figs. 6 and 7 are gaps cut in Black Log and Shade mountains, by a small upper branch stream of the Juniata in southeastern Huntingdon county. The stream traverses a breached antiform of Medusa sandstone, of which these mountains are the nearest remnants. A long narrow valley is opened to the axial Trenton limestone between the two. The gaps are not opposite to each other, and therefore in looking through either gap from the outer country the even crest of the further ridge is seen beyond the axial valley. The gap in Black Log mountain, fig. 6, is located on a small fracture, but in this respect it is unlike most of its fellows.\* The entire region is so strongly characterized by the Medina ridges of the central district. Fig. 8 is in good part an ideal view, based on sketches on the



upper Susquehanna, and designed to present a typical illustration of the more significant features of the region. It shows the even crest-line of a high Medina or Pottsville ridge in the background, retaining the form given to it in the Cretaceous cycle; the even lowlands in the foreground, opened on the weaker Siluro-Devonian rocks in the Tertiary cycle; and the uneven ridges in the middle distance marking the Oriskany and Chemung beds of intermediate hardness that have lost their Cretaceous level and yet have not been reduced to the Tertiary lowland. The Susquehanna flows distinctly below the lowland plain, and the small side streams run in narrow trenches of late Tertiary and Quaternary date.

If this interpretation is accepted, and the Permian mountains are seen to have been once greatly reduced and at a later time worn out, while the ridges of to-day are merely the relief left by

\* Second Geol. Surv. Pa., Report T., 19.

the etching of Tertiary valleys in a Cretaceous baselevelled lowland, then we may well concur with Powell that "mountains cannot remain long as mountains; they are ephemeral topographic forms."<sup>2</sup>

PART TWO. *General conception of the history of a river*

15. *The complete cycle of river life - youth, adolescence, maturity and old age.* The general outline of an ideal river's history may be now considered, preparatory to examining the special history of the rivers of Pennsylvania, as controlled by the geological events just narrated.

Rivers are so long lived and survive with more or less modification so many changes in the altitude and even in the structure of the land, that the best way of entering on their discussion seems to be to examine the development of an ideal river of simple history, and from the general features thus discovered, it may then be possible to unravel the complex sequence of events that leads to the present condition of actual rivers of complicated history.

A river that is established on a new land may be called an original river. It must at first be of the kind known as a consequent river, for it has no ancestor from which to be derived. Examples of simple original rivers may be seen in young plains, of which southern New Jersey furnishes a fair illustration. Examples of essentially original rivers may be seen also in regions of recent and rapid displacement, such as the foot of the broken country of southern Idaho, where the directly consequent character of the drainage leads us to conclude that, if any rivers occupied these regions before their recent deformation, they were so completely extinguished by the newly made slopes that we see nothing of them now.

Once established, an original river advances through its long life, manifesting certain peculiarities of youth, maturity and old age, by which its successive stages of growth may be recognized without much difficulty. For the sake of simplicity, let us suppose the land mass, on which an original river has begun its work, stands perfectly still after its first elevation or deformation, and so remains until the river has completed its task of carrying away all the mass of rocks that lies above its baselevel. This lapse of time will be called a cycle in the life of a river. A complete

<sup>2</sup> *Great Living Mountains*, 874-96.



cycle is a long measure of time in regions of great elevation or of hard rocks, but whether or not any river ever passed through a single cycle of life without interruption we need not now inquire. Our purpose is only to learn what changes it would experience if it did thus develop steadily from infancy to old age without disturbance.

In its infancy, the river drains its basin imperfectly, for it is then embarrassed by the original inequalities of the surface, and lakes collect in all the depressions. At such time, the ratio of evaporation to rainfall is relatively large, and the rate of transported land waste or *talafas*, is small. The channels followed by the streams that compose the river as a whole are narrow and shallow, and their number is small compared to that which will be developed at a later stage. The divides by which the sub-streams are separated are poorly marked, and in level countries are surfaces of considerable area and not, one at all. It is only in the later maturity of a system that the divides are reduced to lines by the consumption of the softer rocks on either side. The difference between constant, *nas* forms and those forms that are due to the action of denuding forces is in a general way so easily recognized, that immaturity and maturity of a drainage system can be readily determined. In the truly infantile drainage

of the great River of the North, the sub-stream areas are so absolutely flat that that water collects on them in wet weather, not having either original structural slope or subsequently developed denuded slope to lead it to the streams. In the almost equally young and unworked western Oregon, the well-marked slopes are as yet hardly channelled by the flow of rain down them, and the depressions among the softer rocks are still undrained, as they have been.

As the river becomes a *nas*, the channels are deepened and the *talafas* increased to considerable extent. If there be a contrast of hardness a low aquia deepening of the down stream part of the channel, while the part next upstream exists above as a cascade or waterfall results, but like the babes of early youth, it is exposed, and continues not a small part of the whole cycle of growth. In the falls on the small mountain streams of a river may last out its maturity, just as here are *young* twigs on the branches of a large tree. With the decreasing of the height, there comes an increase in the number of gulches on the slopes of the channel, the gulches grow to ravines and these

to side valleys, joining the main stream at right angles (La Vie and Margeriel). With their continued development, the number of the system is increased; it is marked by an almost complete obliteration of every part of the original constructed surface, and a new one is formed under the guidance of the stream, so that every point of land that falls finds a way prepared to lead it to a stream and then to the ocean its goal. The lakes of a youthful imperfection have long since disappeared, the waterfalls of a adolescence have been worn back, unless on the still young headwaters. With the increase of the number of sub-streams, ramifying into all parts of the drainage basin, there is a proportionate increase in the surface of the valley slopes, and with this comes an increase in the rate of waste under atmospheric forces, hence it is at maturity that the river receives and carries the greatest load; indeed the increase may be carried so far that the lower trunk stream, of gentle slope in its early maturity, is unable to carry the load brought to it by the upper branches, and therefore resorts to the temporary expansion of its valley, as we saw in a flood plain. The level of the flood plain is sometimes laid up faster than the small sub-streams of the lower course can fill their valleys, and hence they are converted for a brief distance above the mouth into shallow lakes. The growth of the flood plain is so rapid as to carry up the point of junction of tributaries farther and farther from stream, and at last it turning several streams aside from the main stream, sometimes forcing them to follow independent courses to the sea (Long Island). But a thought thus separated from the main trunk, it would be no more rational to regard such stream as an independent river than it would be to regard the branch of an old tree, now fallen to the ground as a tree. If a young tree grows as an independent plant, both are detached to flourish as a single individual, from which they have been separated by an increase of growth and decay.

In the later and quieter end age of a river system, the waste of the land is yielded away by reason of the diminishing slopes of the valley sides, then the headwater streams let ver less detract from the main river, which, thus relieved, turns to its postponed task of carrying as far as it can excess of load to the sea, and gets it rid of in its flood plain, preparatory to sweeping it away. It does this by depositing its load in a new position, and perhaps settling down on a low spot a little to one side of its old line, produces a rapid or a low fall on the lower slope of such an obstruction (Peck). Such a rise may be called locally superimposed.

In early, middle, and mature life, too, as in after years, as that the three divisions of a river, commonly recognized, appear most distinctly; the torrent portion being the still young head-water branches, growing by growing backwards at their sources, the vandy portion proper, where longer time of work has enabled the valley to obtain a greater depth and width, and the lower flood-plain portion, where the temporary deposition of the excess of load is made until the activity of middle age is past.

Maturity seems to be a proper term to apply to this long enduring stage, for as in organic forms, where the term first came into use, it here also signifies the highest development of all functions between a youth of endeavor towards better work and an old age of relinquishment of faintest powers. It is the mature river in which the rainfalls best lead away to the sea, and which carries with it the greatest load of land waste; it is at maturity that the regular descent and steady flow of the river is best developed, being the least delayed in lakes and least overhastened in impetuous falls.


Maturity past, and the power of the river is on the decay. The relief of the land diminishes, for the streams no longer deepen their valleys although the hill tops are degraded; and with the general loss of elevation, there is a failure of runoff to a certain extent, for it is well known that up to certain considerations, the amount of rainfall is proportional to the elevation. A topographic and a hypsometric map of a country for this reason show a marked correspondence. The slopes of the headwaters decrease and the valley sides widen so far that the land waste descends from them slower than before. Later, what with failure of rainfall and decrease of slope, there is perhaps a return to the early imperfect condition of drainage, and the number of side stream and tributaries as branches laid from a dying tree. The flood-plains of maturity are carried down to the sea, and at last the river settles down to an old age of weariness met with gentle flow and light load, little work remaining to be done. The great task that the river entered upon is completed.

10. *Mutual adjustment of river courses*.—In certain structures, chiefly those of more ancient date, on which the streams are at first high above base-level, there is a process of adjustment extremely characteristic of quiet river development, by which the down-hill courses that were chosen in early life, and as we may say unadvisedly and with the heedlessness and little foresight of

youth, are given up for others better fitted for the work of the mature river system. A change of this kind happens when the young stream taking the lowest line for its guide happens to flow on a hard bed at a considerable height above base-level, while its branches or one side or the other have opened channels on softer beds: a part of the main channel may then be deserted by the withdrawal of its upper waters to a lower course by way of a side stream. The change to better adjustment also happens when the initial course of the main stream is much longer than a course that may be offered to its upper portion by the backward growing of an adjacent stream (Lôw, Penck). Sometimes the lateral cutting or planation that characterizes the main trunk of a mature river gives it possession of an adjacent smaller stream whose head is at a higher level (Gibbert). A general account of these processes may be found in Philippon's serviceable "*Etudes sur l'assèchement*" (Leipzig, 1886). This whole matter is of much importance and deserves deliberate examination. It should be remembered that changes in river courses of the kind now referred to are unconnected with any external disturbance of the river basin, and are purely normal spontaneous acts during advancing development. Two examples, pertinent to our special study, will be considered.

Let AB, fig. 9, be a stream whose initial subsequent course led it down the gently sloping axial trough of a syncline. The constructed surface of the syncline is shown by contours. Let the succession of beds to be discovered by erosion be indicated in a section, and in proper position on the several diagrams, but reversed into the horizontal plane, the harder beds being dotted and the base-level standing at 00. Small side streams will soon be developed on the slopes of the syncline, in positions determined by cross-fractures or more often by what we call accident; the action of streams on similar synclines on the outside of the enclosing anticlines will be omitted for the sake of simplicity. In time, the side streams will cut through the harder upper bed M and enter the softer bed N, on which regular, uniform, channels, indicated by hachures, will be extended along the strike, fig. 10 (La Voû and Margerie). Let these be called "subsequent" streams. Consider two side streams of this kind, C and D, heading against each other at E, one joining the main stream lower down the axis of the syncline than the other. The headwaters of C will rob the headwaters of D, because the deepening of the channel

of D is retained by its having to join the main stream at a point where the hard bed in the axis of the fold holds the main channel.



we have likewise. The notch cut by D will then be changed from a water-gap to a wind-gap and the upper portion of B will head out through the notch cut by C, as in fig. 11. As other subsequent headwaters make capture of C, the greater depth to which the lateral valley is cut on the soft rock causes a slow migration of the divides in the abandoned gaps towards the main stream, and before long the upper part of the main stream itself will be diverted from the synclinal axis to follow the more central valley at its side for a distance, fig. 12, until the axis can be rejoined through the gap where the axial portion of the controlling axis was near or at mass level. The upper part of the synclinal trough will then be tracked by undercutting on the slope of the quick-widened channels of the lateral streams, and the hard bed will be worn away in the higher part of the axis before it is



med in the lower part. The location of the successful stream on one or the other side of the syncline may be



from where a hole had been  
 drilled to 1 mile or more  
 and returned to the surface  
 and returned to the surface

\* See the terminology suggested by Powell. Expl. C. d. R. of the West  
 1857. This terminology is applied to the most ordered  
 study of our rivers, by reason of their crossing so many folds, and  
 causing so often for a rapid flow to traverse a fold.

transverse gap, EB, in the next higher hard bed, and there



*Terminology of rivers changed by adjustment.*—A type of terminology is needed for easy reference to the several parts of the streams concerned in such an adjustment. Let AB and CD, fig. 14, be streams of unequal size entering gaps, H and G, in a ridge that lies transverse to their course. CD being larger than AB will deepen its gap faster. Of two subsequent streams, JE and JF, growing on the upstream side of the ridge, JE will have the steeper slope, because it joins the deeper master stream. The divide J, will therefore be drawn toward AB, and if all the conditions concerned compare favorably, JE will at last join AB at F, and lead the upper part, AF, out by the line FECD, fig. 15.




FIG. 15

through the deeper gap, *ie.* We may then say that JE becomes the *divertor* of AF, which is *diverted*, and when the process is completed, by the transfer of the divide from J to the soft rocks, to a stable position, H. In the hard rocks, there will be a short *diverted* stream, HF; while HB is the remaining *beheaded* portion of the original stream, AB, and the water-gap of AB becomes a *wind gap*, H. It is very desirable that geographic exploration should discover examples of the process of adjustment in its several stages. The preparatory stage is easily recognized by the difference in the size of the two main streams, the difference in the depth of their gaps, and the unusual vertical position of the divide, J. The very last stage of transition occurs in the new examples of beheaded streams. For a short time after capture of the diverted stream by the divertor, the new divide will be between F and H, in an unstable position, the duration of this time depending on the energy of the process of capture.

The consequences resulting from rapid incision of this kind by which their recent existence can be detected are: a relatively sudden increase of volume of the diverter and hence a rapid deepening of the course of the diverting stream,  $bb_1$ , and of the diverted,  $Ab$ , near the point of capture, small side streams of these two having unable to keep pace with the change will join their masters in local rapids, which work up stream gradually and fade away (Low, Peck, McGee). The expanded portion,  $ED$ , of the larger stream,  $CD$ , already of fault ~~is~~ ~~is~~ really overcome for a time with the increase of detritus that will be thus conveyed to it at the entrance,  $E$ , of the divertor; ~~where~~ the beheaded stream,  $BB$ , will find itself embarrassed to live up to the habits of its large valley [them]. (Geographic exploration



FIG. 10




FIG. 11




FIG. 12

with these matters in mind offers opportunity for the most attractive of exercises.

18. *Examples of adjustment* - Another case is roughly figured

in the next three diagrams, pgs. 15, 17, 18. Two adjacent synclinal streams, EA and HB, in a transverse master stream, C, and the synclines are of a different form; the surface axis of EA, EA<sub>s</sub>, stands at a much higher baselevel until it nearly reaches the part of the transverse stream; while an axis of the other, HB<sub>s</sub>, descends near baselevel at a considerable distance from the transverse stream. At interval valleys, E and D, are opened on the anticline between the synclines by a process similar to that already described, the surface segment of them will shift towards the stream of fairer slope, that is, to, while the syncline, EA, whose axis is below its hard beds above baselevel; and in time the upper part of the main stream will be withdrawn from this syncline to follow an easier course by crossing to the other, as in fig. 17. If the elevation of the synclinal axis, AEA<sub>s</sub>, take the shape of a long flat area, descending at the farther end into a synclinal lake basin, S, whose outlet is along the arching axis SA<sub>s</sub>, then the mature arrangement of stream courses will lead the lake outlet away from the axis by some gap in the nearer ascending part of the arch where the controlling hard bed fails near to baselevel, as at F, fig. 18,\* and will take it by some subsequent course, F D, across the swland that is opened on the soft beds between the synclines, and carry it into the lower syncline, HB, at D where the hard beds descend below baselevel.

The variety of adjacents following the general principle here indicated is infinite. Changes of greater or less value are introduced in the initial drainage areas, and, after attaining an attack of equilibrium, further change is arrested, or if occurring, is relatively insignificant. It should be noticed that the new stream courses thus chosen are not named by any of the terms now current to express the relation of stream and land history; they are neither consequent, antecedent nor superimposed. The stream is in youth, an original stream, a though no longer young, but its channel is not in all parts strictly consequent on the original contour of land that created it. Streams thus rearranged may therefore be named original streams of mature adjustment.

It should be clearly recognized that the process of adjustment is a very slow one, measured in the extremely long units

\* This figure would be improved if a greater amount of wasting around the margin of the hard bed were indicated in comparison with the preceding figure.

of a river's life. It progresses no faster than the weathering away of the slopes of a divide, and here as a rule weathering is elaborate to say the least, less accelerated by a fortuitous combination of favoring conditions. Among these are high, great altitude if the mass exposed to erosion starts first, and deep channeling of streams by the surface—that is, the advanced stage of drainage development—stands second. The opportunity for the lateral migration of a divide will depend on the inequality of the slopes on its two sides, and here the most important factors are length of the two opposite stream courses from the water parting to the common base-level of the two, and inequality of structure by which one stream may have an easy course and the other a hard one. It is manifest that such these conditions for active shifting of divides are best united in young and high mountain ranges, and hence it is that river adjustments have been found and studied more in the Alps than elsewhere.

10. *Reversal of rivers by elevation and drowning by depression.*

I make no contention that any river in the world ever passed through a simple or interrupted cycle of the orderly kind here described. But by examining many rivers, some young and some old, I do not believe that this portrayal of the idea would be found to be fairly correct if opportunity were offered for its development. The intention of the sketch is simply to prepare the way for the better understanding of our actual rivers of more complicated history.

At the close or at any time during the passage of an entire cycle such as the one just considered, the drainage area of a river system may be bodily elevated. The river is then turned back to a new youth and enters a new cycle of development. This is an extremely common occurrence with rivers, whose life is so long that they commonly outlive the duration of a quiescent stage in the history of the land. Such rivers may be called revived. Examples may be given in which streams are now in their second or third period of revival, the elevations that separate their cycles following so soon that but little work was accomplished in the quiescent intervals.

The antithesis of this is the effect of depression, by which the lower course may be drowned, flooded or fjorded. This change is, if slow, favorable to the development of flood plains in the lower course, but it is not essential to their production. If the change is more rapid, open estuaries are formed, to be transformed to deltas when later on

As a preparation for our subsequent work with regard to the effects of the reversal of a river by *Fig. 31*, let us suppose a new adjustment of its course, by a greater or less extent of a reversal of the new relation of base level to the land.

As previously, *Fig. 32*, Synclinal mountains are most easily explained as results of drainage changes of the kind [Denise, Dec. 21st, 1885]. Streams thus rearranged may be said to be adjusted through elevation or reversal. It is to be hoped that, as our study advances, concise names of brief and appropriate form may replace these paraphrases; but at present it seems advisable to keep the desired idea before the mind by a descriptive phrase, even at the sacrifice of brevity. A significant example may be

Let it be supposed that an originally consequent river system has been so advanced in activity on a surface whose structure is, like that of Henry's mass, composed of closely adjacent anticlinal and synclinal folds with rising and falling axes, and that a series of particularly resistant beds composes the upper members of the folded mass. The master stream, *A*, *Fig. 12*, at maturity still resides where the original folds were lowest, but the side streams have, quite as a matter of course from the axes of the synclinals that they first followed, in accordance with the principles of adjustment, been redirected above. The relief of the surface is moderate, except around the synclinal troughs, where the rising margins of the hard or resistant beds are ridges of more or less prominence. The contour lines in *Figure 12* are drawn on the outcrop side of these ridges. Now suppose a general elevation of the region,

even somewhat above *Fig. 12*. The deepening of the river's master stream will be greatly retarded by reason of its having to cross so many ridges of the hard beds, and the expedient opportunity will be given for readjustment by the growth of some diverting stream, *B*, whose beginning on adjacent softer rocks was already made in the previous cycle. This will separate the main river at some downstream point, and draw it nearly all away from its hard path across the synclinal troughs to an easier path across the lowlands that had been opened on the underlying softer mass, leaving only a small beheaded remnant in the lower course. The final arrangement may be indicated in *Fig. 32*. It should be noted that every capture of branches of the initial main stream made

ments, for with increase of volume the channel is deepened and a




Fig. 6. 6




Fig. 20

Butter & the action of the water in the case of the river away the divides is thereby accelerated. In general it may be said that the larger the stream and the less its elevation above base-level, the less likely is it to be diverted by any other stream at all. It is not impossible however that one stream can offer it a better course to the sea; it may also be said that, as a rule, of two equal streams, the headwaters of the one having a longer or a harder course will be diverted by a branch of the stream on the shorter or easier course. Every case must therefore be examined for itself before the kind of re-arrangement that may be expected or that may have already taken place can be discovered.

21. *Antecedent and superimposed rivers.*—It not infrequently happens that the surface, or even a drainage system is more or less fully dissected, either before or after the formation of a fault. Then, in accordance with the rate of disturbance, and

of the rocks, the streams will be more or less rearranged, some of the larger ones persisting in their courses and cutting their channels down almost as fast as the mass below them is raised and offered to their action. It is manifest that streams of large volume and considerable slope are the ones most likely to persevere in this way, while small streams and large ones of moderate slope may be turned from their former courses to new courses consequent on the new construction and form of the land. Hence, after a disturbance, we may expect to find the smaller streams of the former cycle pretty completely destroyed, while some of the larger ones may still persist, these would then be called antecedent streams in accordance with the nomenclature introduced by Powell.\* A full examination of the development of our rivers will probably give us examples of river systems of all degrees of extinction or persistence at times of disturbance.

Since Powell introduced the idea of antecedent valleys and the Median method of river development, it has found explanation in other regions than the one for which it was first proposed, it has found much acceptance. Low's objection to it does not seem to me to be nearly so well founded as his suggestion of an additional method of river development by means of backward headwater erosion and subsequent capture of other streams, as already described. And yet I cannot help thinking that the explanation of transverse valleys as antecedent courses savors of the Median method of explaining a difficult matter. The case of the Green river, to which Powell first gave this explanation, seems well supported; the examples given by Medico in the Himalayas are as good; but still it does not seem advisable to explain all transverse streams in this way, merely because they are transverse. Perhaps one reason why the explanation has become so popular is that it furnishes an escape from the old catastrophic idea that fractures control the location of valleys, and is at the same time fully accordant with the views of the uniformitarian school that have become current in this half of our century. But when it is remembered that most of the streams of a region are extinguished at the time of mountain growth, that only a few of the larger ones can survive, and that there are other ways in which transverse streams may originate it is evi-

\* Engraving of the Colorado River of the West, 1875, 152-153, 160.

† Hilber, *Pal. Mitth.*, xxxv, 1898, 18.



but the possibility of any given transverse section being made at some or several of the suggested points, would be proved by the actual investigation. It may be difficult to find, but it certainly must be searched for; if not then forth-coming the hypothesis, that the river have the same origin, is a mere appearance. For, if we had a river constant in character, a river would be the hypothetical results of that method of origin, then the burden of proof may be said to lie with those who would maintain that an independent origin would locate the river in so specialized a manner. Even if a river persists, it may be an antecedent course, it may not persist, it may be modified, it may be the various adjustments and rearrangements of the original course, rivers so differently adjusted at the same place, the same time, but it is a more or less different thing, a different adjustment, which we are not yet able to explain. It may be a matter of fact as the Appalachians the courses of the present rivers need not coincide with the courses of the pre-Pennsylvanian rivers, even if we start per- mitted in their courses through the growth of the Pennsylvanian and subsequent rocks, and adjusted to the new conditions, as first buried and unbroken, may have greatly displaced them, in accordance with Law's principle.

When the deeper channeling of a stream discovers an unconformity, the stream is present at a time in the courses that were determined in the overlying mass, they are then called superimposed (Powell), inherited (Shaler), or epigenetic (Hiltholzer). Such streams are particularly liable to run over hard beds to others on which the hard beds are avoided; for the first change of channels, when the unconformable cover was still present, was made without any knowledge of the buried rock structure or of the conditions in which the streams would be involved when they encountered it. The examples of falls produced when streams terrace their flood-plains and run on level surfaces already been referred to as a first step, and the river is now flowing in a new bed, and there may be some of the same thing, the transition stage between the consequent courses, determined by the form of the first channel in which their flow began, and the subsequent courses that will be developed there in the future.

1. Simple, compound, composite and complex rivers.—We

have thus far considered an ideal river. I now seem advisable to introduce a few terms with which to indicate concisely certain well marked peculiarities in the history of actual rivers.

An original river has already been defined as one which first takes possession of a land area, or which replaces a completely extinguished river on a surface of rapid deformation.

A river may be simple, if its drainage area is of practically one kind of structure and of one age; like the rivers of southern New Jersey. Such rivers are generally small. It may be composite when its drainage areas of different ages are united in the basin of a single stream. This is the usual case.

A compound river is one which is of different ages in its different parts; as certain rivers of North Carolina, which have old headwaters rising in the mountains, and young lower courses traversing the coastal plain.

A river is complex when it has entered a second or later cycle of development; the headwaters of a compound river are therefore complex, while the lower course may be simple, as its first cycle of development has been completed. It is not until the second cycle of development has entered that the river has entered

When the study of rivers is thus attempted, its necessary complications may at first seem so great as to render it of no value, but in answer to this I believe that it may be fairly urged that, although complicated, the results are true to nature, and if so, we can have no ground of complaint against them. Moreover, while it is desirable to reduce the study of the development of rivers to its simplest form, in order to make it available for instruction and investigation, it must be remembered that this cannot be done by neglecting to investigate the whole truth in the hope of arriving at the great common sense that simplicity of results is an end in itself, and that the more complicated the process, the more the results will be true to nature.

It is with these points in mind that I have attempted to describe the history of the rivers of Pennsylvania. We find in the Susquehanna, which drains a great area in the central part of the state, an example of a river which is at once composite, compound and highly complex. It is the result of three successive cycles of development, and it is at present in its fourth or fifth degree of complexity, its fourth or fifth cycle of development not yet completed. It is therefore a history of search for the earlier stages of streams which may have been abandoned in the processes of nature's development. It will be

seen earlier, the ~~same~~ *geological history* and *happens* a measure of their previous importance, and to this we may ascribe the difficulty that attends the attempt to decipher a river's history from general maps of its stream area. Nothing but a detailed examination of geological structure and history suffices to detect facts and conditions that are essential to the understanding of the result.

If the postulates that I shall now seem unreasoned and the arguments seem overdrawn, error may at least be avoided by not holding fast to the conclusions that are presented, for they are presented only tentatively. I do not feel by any means absolutely persuaded of the correctness of the results, but at the same time deem them worth going out for discussion. The whole investigation was undertaken as an experiment to see where it might lead, and with the hope that it might lead at least to a serious study of our river problems.

#### PART FOURTH *The development of the rivers of Pennsylvania*

20. *Means of distinguishing between antecedent and adjusted consequent rivers.*—The outline of the geological history of Pennsylvania given above affords means of dividing the *long* progress of the development of our rivers into the several cycles which make up their complete life. We must go far back into the past and imagine ancient streams flowing down from the Archean and towards the paleozoic sea; gaining length by addition to their lower portions as the land grew with the building up of successive mountain ranges; for example, if there were a Cambro-Silurian deformation, a continuation of the Green Mountains into Pennsylvania, we suppose that the pre-existent streams that some number have found their way westward to the new coastline; and from the rate of this mountain growth, it is apparent that any stream that born must have advanced far in their history before the greater Appalachian disturbance began. At the beginning of the latter, as of the former, there must have been streams running from the land into the sea, and at times of temporary elevation of the basin and flats of the coal measures, such streams must have had considerable additions to their lower *it* *is* *may* *is* *may* *growing* Archean highlands or mountains, snow-capped, and drained by glaciers for all we can say to the contrary, descending across the Green Mountains, but, by that time worn to moderate relief in the far advanced stage of its

topographic development, and finally flowing across the enormous lowlands of recent appearance. It was across the lower courses of such rivers that the first mountains were formed, and the first step in our problem consists in deciding if possible whether the streams held their courses after the antecedent fashion, or whether they were thrown into new courses by the growing folds, so that a new drainage-system would be formed. Possibly both conditions prevailed; the larger streams holding their courses undisturbed, and the smaller ones disappearing, to be replaced by others as the slopes of the growing surface should demand. It is not easy to make choice in the matter. To decide that the larger streams persisted and are still to be seen in the greater rivers of to-day, only reversed in direction of flow, is certainly a simple method of treating the problem, but unless some independent reasons are found for this choice, it favors of

Moreover, it is difficult to believe that any streams, even if antecedent and more or less persistent for a time during the mountain growth, could preserve till now their pre-Appa-

the alternations of hard and soft rocks through which they have had to cut, and at all the different altitudes above basins, in which they have stood. A better means of deciding the question will be to admit provisionally the occurrence of a completely

with the slopes of the growing mountains; to study out the changes of stream-courses that would result from later disturbances and from the mutual adjustments of the several members of such a system in the different cycles of its history, and finally to compare the courses thus deduced with those now seen. If there be no accord, either the method is wrong or the streams are not consequent but of some other origin, such as antecedent; if the accord between deduction and fact be well marked, varying only where no definite location can be given to the deduced streams, but agreeing where they can be located more precisely, then it seems to me that the best conclusion is distinctly in favor of the correctness of the deductions. For it is not likely, even if it be possible, that antecedent streams should have accidentally taken, before the mountains were formed, just such locations as would have resulted from the subsequent growth of the mountains and from the complex changes in the initial river courses due to later adjustments. I shall therefore follow the deductive

to find this outline, and attempt to trace out the history of a complete region, and to ascertain the nature of the changes with the growth of the central mountain district.

In doing this, it is first necessary to restore the constructional topography of the region, that is, the form that the surface would have had if it had remained in its original form. This involves certain postulates which must be clearly conceived if any measure of confidence is to be gained in the results based upon them.

1. *Postulates of the argument.*—In the first place, I assume an essential constancy in the thickness of the paleozoic sediments over the entire area in question. This is warranted here because the known variations of thickness are relatively of a second order, and are due to local causes, such as the intense Permian folding. The reasons for maintaining that the whole series had a considerable extension southward of the present margin of the Medina sandstone have already been presented.

In the second place, I shall assume that the dips and folds of the beds now exposed at the surface of the ground may be projected upwards into the air in order to restore the form of the original surface. This is a very delicate question, for it cannot be assumed that the folded slates and sandstones of the Nittany valley, for instance, give any close indication of the original surface. The surface was undoubtedly much more over the district, unworn. But in a general way, the Nittany mass was a high, rugged, and mountainous as well as the Cambrian beds; for our purpose and in view of the moderate relief of the rocks, it is probably safe to assume that wherever the lower rocks are now revealed in antiformal structure, there was a great height and elevation of the original surface, and a corresponding depression of the present surface, which was then a small elevation.

In the third place, I assume that by reconstructing from the completed folds the form which the country would have had if now in the original form, we can picture the form of the surface which it actually passed at the time of initial and progressive folding. The difference between the form of the folds now before us and the form of the original surface is not really a matter of fact, but of degree, and the two are not really so different in general distribution, though and now great as they are.

consideration in our problem. When we remember how accurately water finds its level, it will be clearer that what is needed in the discussion is the location of the regions that were relatively raised and lowered, as we shall then have marked out the general course of the consequent water ways and the trend of the intervening constructional ridges.

Accepting these postulates, it may be said in brief that the outlines of the formation as at present exposed are in effect so many contour lines of the old constructional surface, on which the Permian rivers took their consequent courses. Where the Trenton limestone is now seen, the greatest amount of overlying strata must have been removed; hence the outline of the Trenton formation is our highest contour line. Where the Helderberg limestone appears, there has been a less amount of material removed, hence the Helderberg outcrop is a contour of less elevation. Where the coal beds still are preserved, there has been most wasting, and these beds therefore mark the lowest contour of the early surface. It is manifest that this method assumes that the present outcrops are on a level surface; this is not true, for the ridges through the State rise a thousand feet more or less over the intervening valley lowlands, and yet the existing relief does not count for much in discussing the enormous relief of the Permian surface that must have been measured in tens of thousands of feet at the time of its greatest strength.

*25. Constructional Permian topography and consequent drainage.*

A rough restoration of the early constructional topography is shown in Fig. 21 for the central part of the State, the adjacent areas being the area of the Trenton limestone, indicating the highest ground, or better, the places of greatest elevation, while the Carboniferous area is shaded, indicating the early lowlands. The prevalence of north and southwest trends was then even more pronounced than now. Several of the stronger supports of fact deserve notice, for convenient reference. Thus we have the great Kattunoy or Cumberland highland, 400 on the south-east, marked by the older materials of Cambrian and Archean rocks, falling by the high that may slope to the general lowland troughs of the central district. In this lower ground lay the synclinal troughs of the eastern coal regions, and the more local Broad Top basin, B.F., on the southeast, then better than now preserving the name of basins. Beyond the corrugated area that connected the coal basins rose the great Nittany upland, N,



FIG. 72. Generalized Topographical Map of Pennsylvania.

and its southwest extension in the Bedford range, with the less conspicuous Keshong and beyond, Keshong and beyond the great Allegheny and the names thus suggested are compounded of the local names of to-day and the morphological names of Permian time.

What would be the drainage of such a country? Deductively we are led to believe that it consisted of numerous streams as marked in full lines on the figure, following synclinal axes and some master streams led them across the intervening antoclinal ridges at the lowest points of their crests and away into the open country to the northwest. All the enclosed basins would hold lakes, overflowing at the lowest part of the rim. The general discharge of the whole system would be to the northwest. Here and there we must suppose a channel for the overflow and these well marked features of the great drainage system. The master stream of the region is the great Anthercote river, carrying the overflow of the Anthracite lakes off to the northwest and there perhaps turning along one of the faintly marked streams of the present day, the great Ohio, which was thus continued in its previous location across the Carboniferous marshes. The synclinal streams that entered the Anthercote from the south, the Swatara, S, fig. 21, the Wisconsin, Wo, the Tuscarora, Ma, the Virginia, V, and the Delaware, D, the Wyoming, Wy. One of these, probably the fourth, led the overflow from the Broad Top lake into the Catawba lake on the middle Anthercote river. The Niagary highland formed a strong divide between the central and northwestern rivers, and on its outer slope there must have been a series of streams flowing to the westward and some of these may be regarded as the lower courses of Carboniferous rivers, that once rose in the Archean mountains, now buried by the great Permian range, a rose that is now

20. *The Jura mountains homologous with the Permian Alleghenians.*—However willing one may be to grant the former existence of such a range as the latter, no example of a similar one still in existence would be acceptable as a witness to the possibilities of the past. Therefore we turn for a moment to the Jura mountains and compare them to the Alleghenians on account of the regular series of folds by which the two are characterized. But when the initial topography is long lost in our old mountains, it is still clearly perceptible in the young Jura,





present we look only at the location of the streams, not at their direction of flow. As far as this illustration goes, it gives evidence that the smaller streams at least possess certain peculiarities that are not accounted for by the present topography of the region. It is true that the larger streams were long ago consequent on the Permian folding receives confirmation; but this says nothing as to the origin of the larger rivers, which might at the same time be antecedent.

20. *Departure of the Juniata from the Juniata-Catawissa syncline.*—It may be next noted that the drainage of the Broad Top region does not follow a single syncline to the Allegheno region, as it should have in the initial stage of the subsequent Permian drainage, but soon turns aside from the syncline in which it starts and runs across country to the Susquehanna. It is true that in its upper course the Juniata departs from the Broad Top region by one of the two synclines that were indicated as the probable one of discharge of the ancient Broad Top lake in our restoration of the reconstructed topography of the State; there does not appear to be any significant difference between the summit altitudes of the Tuscarora-Mahanoy and the Juniata-Catawissa synclines and hence the choice must have been made for reasons that cannot be detected; or it may be that the syncline lying there to the northwest was raised last, and for this reason was taken as the one of overflow. The beginning of the river is therefore not discordant with the hypothesis of consequent drainage, but the southward departure from the Catawissa syncline at Lewisburg remains to be explained. It seems to me that some reason for the departure may be found by likening it to the case already given in figs. 16-18. The several synclines with which the Juniata is concerned have precisely the relative positions that are there indicated. The Juniata-Catawissa syncline has parallel sides for many miles about its middle, and must have long maintained the level Juniata well above baselevel over all this distance; the progress of cutting down a channel through all the hard Carboniferous sandstones for a great distance along the axis must have been exceedingly slow. But the synclines next south, the Tuscarora-Mahanoy and the Wiconisco, plunge to the northeast more rapidly, so the rapid

divergence of their margins demonstrates, and must for this reason have earned the hard sandstone brow baselevel in a shorter distance and on a steeper slope than in the Onawissa syncline. The further southwestward extension of the Piceano sandstone ridges in the southern than in the northern ~~direction~~ gives further illustration of this peculiarly of form. Lateral capture of the Juniata by a branch of the initial Tazewell, and of the latter by a branch of the Wisconsin therefore seems possible, and the accordance of the facts with so highly specialized an arrangement is certainly again indicative of the correctness of the hypothesis of consequent drainage, and this time in a larger stream than before. At first sight, it appears that an easier lateral capture might have been made by some of the streams flowing from the outer slope of the Nittany highland; but this becomes improbable when it is perceived that the heavy ~~Massachusetts~~ ~~beds~~ ~~of the~~ ~~Carboniferous~~ ~~series~~ ~~in the~~ ~~Seven~~ ~~Mountains~~ ~~area~~ ~~are~~ ~~not~~ ~~so~~ ~~highly~~ ~~developed~~ ~~as~~ ~~the~~ ~~repeated~~ ~~arches~~ ~~of~~ ~~the~~ ~~Carboniferous~~ ~~beds~~ ~~in~~ ~~the~~ ~~many~~ ~~high~~ ~~folds~~ ~~of~~ ~~the~~ ~~Seven~~ ~~Mountains~~. Again, as far as present appearances go, we can give no sufficient reason to explain why possession of the headwaters of the Juniata was not gained by the subsequent stream of its own, such as to, fig. 18, instead of by a side-stream of the river in the neighboring syncline, but it may be admitted, on the other hand, that as far as we can estimate the chances for conquest, there was nothing distinctly in favor of one or the other of the side-streams concerned; and as long as the problem is solved indifferently in favor of one or the other, we may accept the lead of the facts and say that some control not now apparent determined that the diversion should be, as drawn, through D and not through G. The detailed location of the Juniata in its middle course below Lewistown will be considered in a later section.

3. *Abolition of the Broad Top basin by the Juniata headwaters.*—Another highly characteristic change that the Juniata has suffered is revealed by examining the adjustments that would have taken place in the general topography of the Broad Top district during the Perm-Triassic cycle of erosion. When the basin, BT, fig. 22, was first outlined, centripetal streams descended its slopes from all sides and their waters accumulated as a lake in the center, overflowing to the east into the subterranean ~~basin~~ ~~of~~ ~~the~~ ~~Juniata~~ ~~system~~ ~~along~~ ~~one~~ ~~of~~ ~~its~~ ~~larger~~ ~~basins~~, and thence escaping northeast. In due time, the



on only the weakest rocks, but after this, the stream had grown to a good-sized river, farther rising of the land, probably in the time of the Jurassic elevation, allowed the river to sink its channel to a greater depth, and in doing so, it encountered the hard Medusa anticline of Jack's mountain; here it has never penetrated, because, as we may suppose, there has been no stream able to divert the course of so large a river from its growing of a single hard anticline.

The doubt that one must feel as to the possibility of the processes just outlined arises, if I may gauge it by my own feeling, rather from incredulity than from direct objections. It seems incredible that the waste of the valley slopes should allow the backward growth of N at such a rate as to enable it to capture the heads of C, Tn, k, and so on, before they had cut their heads down close enough to the baselevel of the time to be safe from capture. But it is difficult to urge explicit objections against the process or to show its quantitative inefficiency. It must be remembered that when these adjustments were going on, the region was one of great altitude, its rocks then had the same strong contrast of strength and weakness that are so apparent in the present relief of the surface and the streams concerned were of moderate size; has been now, for at the time, the Tyrone, Frankstown and Bedford head branches of the Juniata had not acquired drainage west of the great N-tiny-Bedford an axial axis, but were supplied only by the rainfall on its eastern slope (see section 39, and all these conditions conspired to favor the adjustment. Finally, while apparently extraordinary and difficult

1 demonstration, the explanation if applicable at all certainly gives no such correlation to a number of peculiar and specific stream courses in the upper Jamuna district that are meaningless under any other theory that has come to my notice. It is chiefly for this reason that I am inclined to accept the explanation.

11. *Reversal of larger rivers to southeast courses*—Our large rivers at present flow to the southeast, not to the northwest. It is difficult to find any precise date for this reversal of flow from the former hypothetical direction, but it may be suggested that it occurred about the time of the Triassic depression of the Newark belt. We have been persuaded that much time elapsed between the Permian folding and the Newark depression, even under the most liberal allowance for a Permian or even a late Newark belt—hence when the depression began, the rivers must

have had but moderate northwestward declivity. The upheaval and submergence of the broad Newark belt may at this time have broken the continuity of the streams that once flowed across it. The headwater streams from the ancient Appalachian country maintained their courses to the depression, the lower portions of the rivers may also have gone on as before; but the middle courses were perhaps turned from the central part of the state back of the Newark belt. No change of altitude gives a fitting cause of the southeastward flow of our rivers as this. The only test that I have been able to devise for the suggestion is one that is derived from the relation that exists between the location of the Newark belt along the Atlantic slope and the course of the neighboring transverse rivers. In Pennsylvania, where the belt reaches somewhat beyond the northwestern margin of the crystalline rocks in South Mountain, the streams are reversed, as above stated; but in the Carolinas where the Newark belt lies far to the east of the boundary between the Cambrian and crystalline rocks, the Tennessee streams preserve in what we suppose to have been their original direction of flow. This may be interpreted as meaning that in the latter region, the Newark depression was not felt distinctly enough, if at all, within the Appalachian belt to reverse the flow of the streams; while in the former region, it was nearer to these streams and determined a change in their courses. The original Anthracite river ran to the northwest, but its middle course was afterwards turned to the southeast.

I am free to allow that this has the appearance of heap-hypothesis on hypothesis; but in no other way does the analysis of the history of our streams seem possible, and the success of the experiment can be judged only after making it. At the same time, I am constrained to admit that such is to my own view the least satisfactory of the suggestions here presented. It may be said that there seems to be too sufficient exclusion of other hypotheses. For example, it must not be overlooked that, if the Anthracite river ran southeast during Newark deposition, the formation of the Newark northwestward topoclinal by the Jurassic tilting would have had a tendency to turn the river once again to its northwest flow. But as the drainage of the region is still southeastward, I am tempted to think that the Jurassic tilting was not here strong enough to reverse the flow of so strong and mature a river as the Anthracite had by that time.

come to be ; and that the elevation that accompanied the tilting was not so powerful in reversing the river to a northwest course as the previous depression of the Newark basin had been in turning it to the southeast. If the Appalachians did continue to draw to the southeast, it may be added that the down-cutting of its upper branches was greatly retarded by the decrease of slope in its lower course when the monocline was formed.

The only other method of reversing the original northwestward flow of the streams that I have imagined is by capture of the headwaters by Atlantic rivers. This seems to me less effective than the method just considered, but they are not mutually exclusive and the actual result may be the sum of the two processes. The outcome of the idea is as follows. The long continued supply of sedimentary material from the Archean land on the southeast implies that it was so continually elevated. But here came a time when there is no record of farther supply of material, and when we may therefore suppose the elevation was no longer maintained. From that time onward, the Archean range must have denuded away, what with the encroachment of the Atlantic on its eastern shore and the general action of denuding forces on its surface. The Newark depression was as effective and in the same way, as has been stated above, and for a moderate distance westward of the depressed belt, the former direction of the streams must certainly have been reversed ; but the question remains whether the reversal extended as far as the Wyoming basin, and whether the subsequent formation of the Newark monocline did not undo the effect of the Newark depression. It is manifest that as far as our limited knowledge goes, it is impossible to estimate these matters quantitatively, and hence the importance of looking for additional processes that may supplement the effect of the Newark depression and counteract the effect of the Newark uplift in changing the course of the rivers. Let it be supposed for the moment that at the end of the Jurassic uplift by which the Newark monocline was formed, the divide between the Ohio and the Atlantic drainage lay about the middle of the Newark belt. There was a long gentle descent westward from the watershed and a shorter and hence steeper descent eastward. Under such conditions, the divide must have been pushed westward, and as long as the rocks were so exposed as to open areas of weak sediments on which capture by the Atlantic streams could go on with relative rapidity, the westward migration of the

divide would be important. For this reason, it might be carried from the Newark bed as far as the present Alleghany front, beyond which further pushing would be slow, on account of the broad stretch of country there covered by hard horizontal beds.

The end of this is that, under any of the circumstances here outlined, there would be early in the Jurassic-Cretaceous cycle a distinct tendency to a westward migration of the Atlantic-Ohio divide; it is the consequences of this that have now to be examined.

32. *Capture of the Anthracite headwaters by the growing Susquehanna*.—Throughout the Perm-Triassic period of denudation, a great work was done in wearing down the original Alleghenian Anticline of hard sandstone, and broad lowlands were opened on the softer rocks beneath. Little remnant of the early constructional topography remained when the period of Newark depression was brought to a close; and all the westward headwater streams of the region were gushing at the divide, seeking to develop the most perfect arrangement of waterways. Several adjustments have taken place, and the larger streams have been reversed in the direction of their flow; but a more serious problem is found in the disappearance of the original master stream, the great Anthracite river, which must have at first led away the water from all the lateral synclinal streams. Being a large river, it could not have been easily diverted from its course, unless it was greatly retarded in cutting down its channel by the presence of many beds of hard rocks on its way. The following considerations may perhaps throw some light on this obscure point.

It may be assumed that the whole group of mountains formed by the Permian deformation had been reduced to a moderate relief when the Newark depression was stopped by the Jurassic elevation. The harder ribs of rock doubtless remained as ridges projecting above the intervening lowlands, but the strength of relief that had been given by the constructional forces had been lost. The general distribution of residual elevations then remaining unobscured is indicated in fig. 25, in which the Crystalline, the Medina, and the two Carboniferous sandstone ridges are denoted by appropriate symbols. In restoring this phase of the surface form, when the country stood lower than now, I have reduced the antiforms from their present outlines and increased the synclines, the change of area being made





greatest where the dips are least, and hence most apparent at the ends of the plunging anticlines and synclines. Some of the Medina anticlines of Perry and Juniata counties are not indicated because they were not then discovered. The country between the residual ridges of Jurassic time was chiefly Cambrian limestone and Siluro-Devonian shales and soft sandstones. The moderate ridges developed on the Onondaga and Chemung sandstones are not represented. The drainage of low stage retained the original courses of the streams, except for the adjustments that have been described, but the great Anthracite river is drawn as if it had been controlled by the Newark depression and reversed in the direction of its flow, so that its former upper course on the Cambrian rocks was replaced by a superimposed Newark lower course. Fig. 15 therefore represents the streams for the most part still following near their synclinal axes, although departing from them where they have to enter a synclinal cover-mountain ridge; the headwaters of the Juniata avoid the mass of hard sandstones discovered in the bottom of Mill Brook Gap lake, and flow around them to the north, and then by a cross-country course to the Wernicke synclinal, as already described in detail. Several streams come from the northeast, entering the Anthracite district after the fashion generalized in fig. 13. Three of the many streams that were developed on the great Antietam slope are shown, with their direction of flow reversed; these are marked S<sub>1</sub>, L, and D, and are intended to represent the ancestors of the existing Susquehanna, Lehigh and Delaware. We have now to examine the opportunities offered to these small streams to increase their drainage areas.

The Jurassic elevation, by which the Newark depression was stopped, restored to activity all the streams that had in the previous cycle sought and found a course close to baselevel. They now all set to work again deepening their channels. But in this restoration of lost activity with reference to a new base-level, there came the best possible chance for numerous re-arrangements of drainage areas by natural adjustment into which we must inquire.

I have already illustrated what seems to me to be the type of the conditions involved at this time in figs. 19 and 20. The master stream, A, traversing the synclines, corresponds to the reversed Anthracite river; the lowlands at the top are those that have been opened out on the Siluro-Devonian beds of the

the actual Susquehanna, the line between the Potomac and the Maryland basins. The actual stream, I think, a great drainage area in these lowlands, corresponds to the embryo of the present Susquehanna, fig. 20, this having been itself once a branch on the north side of the Swatara synclinal stream, fig. 21, from which it was first turned by the anticline to the westward of the Newark depression; but it is located a little farther west than the actual Susquehanna, so as to avoid the two synclinal core mountains of Pocono sandstone that the Susquehanna now traverses, for reasons to be stated below (section 25). This stream had to cross only one bed of hard rock, the outer wall of Medina sandstone, between the broad inner low lands of the relatively weak Siluro-Devonian rocks and the great valley low lands on the still weaker Cambrian limestones. Step by step it must have passed its headwater divide northward, and from time to time it would have thus captured a subsequent stream, that crossed the divide southward, and entered a Carboniferous syncline by one of the lateral gaps already described. With every such capture, the power of the growing stream to capture others was increased. Fig. 19 represents a stage after the streams in the Swatara and Wisconsin synclines (the latter then having gained the Juniata) had been turned aside on their way to the Carboniferous basin. On the other hand, the Anthracite river, rising somewhere on the plains north of the Wyoming syncline and pursuing an irregular course from one coal basin to another, found an extremely difficult task in cutting down its channel across the numerous hard beds of the Carboniferous sandstones, so often repeated in the rolling folds of the coal fields. It is also important to remember that an aid to other basins concerned in the diversion of the upper Anthracite is found in the increase of slope that its lower course suffered in crossing the coal beds. If that area took any part in the deformation that produced the Newark monocline—whichever theory prove true in regard to the origin of the southerlyward flow of the rivers—for loss of slope in the middle course, where the river had to cross the reefs of hard sandstone, would have been very effective in delaying the time allowed for the diversion of the headwaters.

The question is, therefore, whether the retardation in cutting here experienced by the Anthracite was sufficient to allow the capture of its headwaters by the Susquehanna. There can be little doubt as to the correct quality of the process, but

would be sufficient, and that matter. In the absence of any means of testing its sufficiency as the result of taking as the basis is not the correspondence between theory and fact close enough to prove the correctness of the deduction?

*Present outward drainage of the anthracite basin.*—The Lehigh, like the Susquehanna, made an attempt to capture the headwaters of adjacent streams, but failed to acquire much territory from the Anthracite because the Carboniferous sandstones spread out between the two in a broad plateau of hard rocks, a formidable barrier to conquest. The Delaware tried that its upper branches drain is, I think, the conquest of a later cycle of growth. The Delaware had little success, except as against certain eastern synclinal branches of the Anthracite, for the same reason. The ancestor of the Swatara of to-day made little progress against the Anthracite because its first attack was against the repeated Carboniferous sandstones in the Swatara syncline. Consequently it did not find a favorable opportunity for conquest, and thus grew to be the master river—the Susquehanna of to-day. The head of the Anthracite was carried away by this captor, and its beheaded lower portion merged in the Susquehanna. The Allegheny, a tributary formerly of the Lehigh, was carried away as well, being attracted to the surrounding rivers. As the broad Devonian lowlands were opened around the coal basins, especially on the north and west, the streams that formerly flowed into the basins were gradually inverted and flowed out of them, as they still do. The extent of the inversion seems to be in a general way proportionate to its opportunity. The most considerable conquests were made in the upper basins, where the Catawissa and New Hope streams of to-day drain many square miles of wide valleys opened on the Manch Chunk red shale between the Pocono and Pottsville sandstone ridges; the ancient middle waters of the Anthracite here have been diverted to the Susquehanna. In the lower basins the northern coal basins were degraded very slowly after the upper Anthracite had been diverted. The Schuylkill as the modern representative of the Anthracite retains only certain streams south of a medial divide between New Hope and Blue mountains. The only considerable part of the old Anthracite river that still retains a course along the axis of a synclinal trough seems to be that part which flows in the Wyoming basin; none of the many other

val basins are now occupied by the large stream that originally followed them. The reason for this is manifest if to be found in the fact that the West Branch has not been able to do it, if its head ramifications are even now below baselevel, and hence have never yet acted to throw the river from its axial course. Indeed, during the early cycles of denudation, this basin must have been changed from a deep lake to a lacustrine plain by the accumulation in it of waste from the surrounding high lands, and for a time the stream that entered it may have flowed in meandering courses across the ancient subalpine surface; the lacustrine and alluvial condition may have been temporarily revived at the time of the Jurassic elevation. It is perhaps an overstatement from a course thus locally superimposed that we may come to regard the deflection of the river at Natcoke from the axis of the syncline to a narrow shale valley on its northern side, before turning south again and leaving the basin altogether. But like certain other suggestions, this can only be regarded as an open hypothesis, to be tested by some better method of river analysis than we now possess; like several of the other explanations here offered, it is presented more as a possibility than as dismissed than as a conclusion to be accepted.

I believe that it was during the earlier part of the great Jurassic retrogressive cycle of denudation that the Susquehanna took the course she musters across of the central district of the state. For the rest of the cycle, it was occupied in carrying off the waste and reducing the surface to a well-washed baselevel lowland that characterized the end of Cretaceous time. From an active youth of conquest, the Susquehanna advanced into an old age of established boundaries; and in later times, its area of drainage does not seem to have been greatly altered from that so long ago defined, except perhaps in the districts drained by the West and North Branch headwaters.

14. *Homologies of the Susquehanna and Juniata*.—Looking at the change from the Allegheny to the Susquehanna in a broad way, one may perceive that it is an effect of the same order as the peripheral changes of the broad Topographic Age, illustrated in Figures 22, 23 and 24. Another example of a similar change is seen in the later stages of the Juniata above Lewisburg and its north-west continuation in Augwick creek, from their original axial location when they formed the initial broad Topographic outlet. They have departed from the axis of their syncline to

the softer beds on its southern side; *FE* of *fig. 17* has been diverted to *ED* of *fig. 19*.

All of these examples are truly only special cases of the one already described in which the Juniata left its original syncline for others to the south. The general case may be stated in a few words. A stream flowing along a syncline of hard beds (Carboniferous sandstones) develops side streams which breach the adjacent anticlines and open lowlands in the underlying softer beds (Devonian and Silurian). On these lowlands, the headwaters of side streams from other synclines are encountered and a contest ensues as to possession of the drainage territory. The divides are pushed away from those headwaters whose lower course leads them over the fewest hard barriers, the conquest goes on until the upper course of the original main stream is diverted to a new and easier path than the one it chose in its youth in obedience to the first information of the region. Thus the Juniata now avoids the center and once deepest part of the old Wood Top lake, because in the general progress of erosion, lowlands on soft Devonian beds were opened up around the edge of the great mass of sandstones that held the lake; the original drainage across the lake, from its western outlet to its outlet just south of the Jack's mountain anticline, has now taken an easier path along the Devonian beds to the west of the old lake basin, and is seen in the Little Juniata, flowing along the outer side of Terrace mountain and reaching the northern anticline just where Terrace mountain joins Sublingia. It then crosses Jack's mountain at a point where the hard Menara sandstones of the mountain were still buried at the time of the carving of this channel. In the same way, the drainage of the Susquehanna basin through which the main lake discharged eastward, is now not along the axis of the Juniata-Gettysburg syncline, but on the softer beds along the side of it; and along the same line, the drainage of the Potomac that was provided for it by our first folds, namely, via the Incarnata and Wisconsin synclines, as already described. The much broader drainage from the Appalachians to the Susquehanna was only another form of the same process. Taking a transverse view of the whole system of central folds, it is perceived that four axes descend into the Antietam district from the east and rise westward to the surface. It is as if the whole region had received a transverse depression, and the transverse axis of depression thus formed defined the initial course of the first major stream.

But the master stream deserted its original course on the transverse axis of depression because a lateral course across lowlands on softer beds was opened by its side streams. And in the contest of these lowlands with an external stream, the Susquehanna, the upper portion of the An bracts was diverted from the hard rocks that had appeared on the transverse axis. The distance of diversion from the axial to the lateral course in this case was great because of the gentle quality of the transverse folding, or, better said, because of the gentle dips of the axes of the longitudinal folds. This appearance of systematic re-arrangement in the several river courses where none was expected is to my mind a strong argument in favor of the original & consequent location of the rivers and their later mutual adjustment. It may perhaps be observed that antecedent streams might imitate one another roughly in the attitude that they project only those with regard to folds subsequently formed, but no reason has been suggested for the imitation being carried to so remarkable and definite a degree as did here and there.

35. *Superimposition of the Susquehanna on two synclinal ridges.* There is however one extremely venturesome postulate that may have been already noted as such by the reader, unless it can be reasonably accounted for and shown to be a natural result of the long sequence of changes here considered, it will seriously militate against the validity of the whole argument. The present course of the middle Susquehanna leads it through the apical curves of two Pocono synclinal ridges, which were disregarded in the statement given above. It was then assumed that the embryonic Susquehanna gained possession of the soft rock as low level drainage by growing out a course to the west some miles; for it is not to be thought of as any conquest of the headwaters of the An bracts river could have been made by the Susquehanna if it had had to grow out the existing four traverses of the Pocono strata before securing its drainage of the lowlands above them. If a backward progress of the Susquehanna could not in that case have been nearly fast enough to reach the Anticline before the latter had sunk its summit to a safe depth. It is therefore important to justify this assumption as to the more westerly behavior of the embryonic Susquehanna; and afterwards to explain how it should have since then been transferred to its present course. A short cut through all this maze even at present is open to those who adopt

in the beginning the theory that the Susquehanna was an antecedent river; but as I have said at the outset of this inquiry it seems to me that such a method is not free from assumption, even though simpler than the one here adopted, and that it has the demerit of not considering all the various details that follow the examination of consequent and adjusted courses.

The sufficient reason for the assumption that the embryonic Susquehanna lay farther west than the present one in the horizon of the Pocono syncline is simply that, in the absence of any antecedent stream—it must have been there. The whole explanation of the development of the Susco-Deo as a lowland between the Pocono and Medina ridges appears simply on their being weathered and where the rocks are weak enough to waste faster than the enclosing harder ranges through which the streams escape. In this process, the streams exercise no control whatever over the direction in which their headwaters shall grow; they leave this entirely to the structure of the district that they drain. It thus appears that, under the postulate as to the initial location of the Susquehanna as one of the many streams descending the great slope of the Keweenaw (Catharland) highland into the Western syncline its course being reversed from eastward to westward by the Newark depression, we are required to suppose that its headwaters (northward) grew at the time of the Jurassic elevation must have been on the sub-Cretaceous beds, so as to avoid the harder rocks on either side. Many streams competed for the distinction of becoming the master, and that one gained its pre-eminence whose initial location gave it the best subsequent opportunity. It remains then to consider the means by which the course of the embryonic Susquehanna may have been subsequently changed from the lowlands to the two Pocono synclines that it now traverses. Some departure from its early location may have been due to eastward planation in its advanced age, when it had large volume and gentle slope and was therefore scouring and cutting laterally in its lower course. This may have had some share in the result, but there is another process that seems to me more effective.

In the latter part of the Jura-Cretaceous cycle, the whole country hereabout suffered a general rate depression by which the Atlantic transgression many miles inland from its former shoreline, across the low areas of erosion that had been developed on the inland belt. Such a depression must have had a distinct effect



on the lower courses of the larger rivers, which having already cut their channels down close to base-levels and opened their valleys wide on the softer rocks, were then "estimated," or at least so far checked as to build wide flood plains over their lower stretches. Indeed, the flood-plains may have been begun at an earlier date, and have been confirmed and extended in the later time of depression. Is it possible that in the latest stage of the process, the almost base-levelled remnants of Blue mountain and the Pocono ridges could have been buried under the flood plain in the neighborhood of the river?

If this be admitted, it is then natural for the river to depart from the line of its buried channel and cross the buried ridges on which it might settle down as a superimposed river in the next cycle of elevation. It is difficult to decide such general questions as these, and it may be difficult for the reader to gain much confidence in the admissibility of the processes suggested; but there are certain features in the side streams of the Susquehanna that lend some color of probability to the explanation as offered.

Admit, for the moment, that the aged Susquehanna, in the later part of the Jura-Cretaceous cycle, did change its channel somewhat by cutting to one side, or by planation, as it is called. Admit, also, that in the natural progress of its growth it had laid a broad flood plain over the Siluro-Devonian low lands, and that the depth of this deposit was increased by the formation of an estuary delta upon it when the country sank at the time of the final Cretaceous transgression of the sea. It is manifest that one of the consequences of all this might be the peculiar course of the river that is to be explained namely its superimposition on the two Pocono synclinal ridges in the next cycle of its history, after the Tertiary elevation had given it opportunity to re-discover them. It remains to inquire what other consequences would follow from the same conditions, and from these to derive tests of the hypothesis.

1st. *Evidence of superimposition.*—One of the peculiarities of flood-plain rivers is that the lateral streams shift their points of union with the main stream farther and farther down the valley, as indicated in the case of the Po. If the Susquehanna were heavily flood-plained at the close of the Jura-Cretaceous cycle, some of its tributaries should, manifest a good deal of kind of deflection from their structural courses along the strike of the rocks. Some



antecedent river; and yet there is no reason to think that it could have been brought into its present special position by any process of shifting divides. The processes that have been suggested to account for its special location, as departing slightly from a position due to slow adjustments following an ancient consequent origin, call for the occurrence of certain additional peculiarities in the courses of its tributary streams, entirely unforeseen and unlooked for at this point in the inquiry as reached; and in looking at the map to see if they occur, they are found with perfect distinctness. The hypothesis of superimposition may therefore be regarded as having advanced beyond the stage of mere suggestion and as having gained some degree of confirmation from the correlations that it detects and explains. It only remains to ask if these correlations might have originated in any other way, and if the answer to this is in the negative, the case may be looked upon as having a fair measure of evidence in its favor. The remaining question may be taken up at once as the first point to be examined in the Tertiary cycle of development.

37. *Events of the Tertiary cycle.*—The elevations given to the region by which Cretaceous baseleveling was terminated, and which I have called the early Tertiary elevation, offered opportunity for the streams to deepen their channels once more. In doing so, certain adjustments of moderate amount occurred which will be soon examined. As time went on, much denudation was effected, but no widespread baseleveling was reached, for the Cretaceous crest lines of the hard sandstone ridges still exist. The Tertiary cycle was an incision one. At its close, low lands had been opened out on the weaker rocks between the hard beds. Is it not possible that the flood planing of the Susquehanna and the down-stream deflection of its branches took place in the closing stages of this cycle, instead of at the end of the previous cycle? If so, the deflection might appear on the branches, but the main river would not be transferred to the Pocono ridges. This question may be safely answered in the negative, for the Tertiary upland is by no means well enough baselevelled to permit such an event. The beds of intermediate resistance, the Onondaga and certain Cambrian sandstones, had not been worn down to baselevel at the close of the Tertiary cycle; they had indeed lost much of the height that they possessed at the close of the previous cycle, but they had not been reduced as low as the softer beds on either side. They were only reduced to ridges of

moderate but unequal height over the general plain of the Sea. The main low country, without great strength of relief but quite strong enough to cause obedience from the streams along side of them. And yet near Swan's Grove, for example, in Sawyer county, Peter's and Middle creeks depart most distinctly from the strike of the local rocks as they near the Desquehanna, and traverse certain well-marked ridges on their way to the main river. Such aberrant streams cannot be regarded as superimposed at the close of the Tertiary or Quaternary cycle; they cannot be explained by any process of spontaneous adjustment yet described, nor can they be regarded as vastly ancient streams of antecedent courses; I am therefore much tempted to consider them as of superimposed origin, inserting their present courses from the flood-plain cover of the Desquehanna in the latest stage of the Tertiary or Quaternary cycle. With this tentative conclusion on mind as to the final events of Tertiary or Quaternary time, we may take up once more our consideration of the work of the Tertiary cycle.

The chief work of the Tertiary cycle was merely the opening of the valley winds; the opportunity for river adjustment occurred except on a small scale. The most evident cases of adjustment have resulted in the change of water-gaps into wind-gaps, of which several examples can be given, the one best known being the Delaware wind-gap between the Lohigh and Delaware water-gaps in Blue mountain. The wind-gap marks the unfinished notch of some stream that once crossed the ridge here and whose sandwaters have since then been diverted, probably to the Lohigh. The difficulty in the case is not at all how the stream that once flowed here was diverted, but how a stream that could be diverted in the Tertiary cycle could have escaped diversion at some earlier date. The relative rarity of wind-gaps indicates that nearly all of the initial lateral streams, which may have crossed the ridges at an early epoch in the history of the rivers, have been inboarded at some cycle earlier than the Tertiary and their gaps thereafter obliterated. Why the Delaware wind-gap stream should have endured into a later cycle does not at present appear. Other wind-gaps of apparently similar origin may be found in Blue mountain west of the Schuylkill and east of the Desquehanna. It is noteworthy that if any wind streams still persevere in their gaps across a hard ridge, they are not very close to any large river-gap; hence it is only at the very headwaters of Conodogwinit creek, in the

northern part of Franklin county, that any water is still drawn from the back of Blue Mountain. Again, three main stream gaps do not lie between large river-gaps and wind-gaps, but wind-gaps lie between the gaps of large rivers and those of small streams that are not yet diverted. Exact observation of this is found on the "Piedmont sheet" of the colored maps issued by the United States Geological Survey. The sheet covers part of Maryland and West Virginia, near where the North Branch of the Potomac comes out of the plateau and crosses New Creek mountain. Eleven miles south of the Potomac gap there is a deep wind-gap; but further on, at twenty, twenty-five and twenty-seven miles from the river-gap are three fine water-gaps occupied by small streams. This example vividly shows how many important points in the history of our rivers will be missed when the country is properly portrayed on colored maps.

A few lines may be given to the general sequence of gaps in Blue Mountain in Pennsylvanian. When the initial consequent drainage was established many streams must have been located on the northward slope of the great Cumberland high land, (fig. 21), they must have gullied the slope to great depths and carried away great volumes of the weak Cambrian beds that lay deep within the hard outer casings of the mass. Minor adjustments served to diminish the number of these streams, but the more effective cause of their present rarity lay in the nature and location of certain of them to become large streams; the smaller ones were generally beheaded by these. The only examples of streams that still cross this ridge with their initial Permian direction of flow to the north-west are found in two southern branches of Tuscarora creek at the southern point of Juniata county; and these survive because of their obscure location among the many Medina ridges of that district, where they were not easily accessible to capture by other streams.

8. *Tertiary adjustment of the Juniata on the Medina anticline.* The lower course of the Juniata presents several examples of adjustment referable to the last part of the Juratrias cycle and to the Tertiary cycle. The explanation offered for the escape of this river from its initial syncline did not show any reason for its peculiar position with respect to the several Medina anticlines that it now borders, because at the time when it was still normal country to the Wisconsin syncline, the hard Medina heads of these anticlines were not discovered. It is therefore

hardly to be caught at the junction of the Juniata in the Narrows now Lewistown between Blue Ridge and Shade mountains and its avoidance of Tuscarora mountain could have been defined at that early date. But all these Medina anticlines rise more or less above the Cretaceous basalt, and must have had some effect on the point on taken by the river above the mouth of Cretaceous when the channel sank upon them. Blue Ridge and Black Log anticlines rise highest. The first head-on of the cross-country stream that led the early Juniata away from its original syncline probably traversed the Blue Ridge and Black Log anticlines while they were yet buried, but its channel-cutting was much retarded on encountering them, and some branch stream working around from the lower side of the obstructions may have diverted the river to an easier path. The only path of the kind is the narrow one between the overlapping and faces of Blue Ridge and Shade mountains, and there the Juniata now flows.

If an chief elevation should occur in the future, it might happen that the slow deepening of the channel in the hard Medina beds which now floor the Narrows would allow Middle creek of Snyder county to tap the Juniata at Lewistown and lead it by direct course past Middletown to the Susquehanna; then it would return to the path of its youth.

The location of the Juniata at the end of Tuscarora mountain is again so definite that it can hardly be referred to a time when the mountain had not been revealed. The most likely position of the original cross-country stream which brought the Juniata into the syncline was somewhere on the line of the existing depression, and assuming it to have been there, we must question how it has been displaced. The process seems to have been of the same kind as that just given; the retardation of channel-cutting on the late Cretaceous cycles, when the Medina beds of Tuscarora anticline were uncovered, allowed a branch from the lower part of the river to work around the end of the mountain and lead the river out that way. The occurrence of a shallow depression across the summit of the otherwise remarkably even crest of Tuscarora mountain suggests that this diversion was not hardly accomplished until shortly after the Tertiary elevation of the country; but at whatever date the adjustment occurred, it is natural that it should pass around the eastern end of the mountain and not around the western end, where the course would have been much longer, and therefore not successfully to be taken by a diverting stream.

While the quality of these processes appears satisfactory, I am not satisfied as to the sufficiency of their quantity. If diversion was successfully practiced at the crossing of the Tuscarora anticline, why not also at the crossing of Jack's mountain anticline, on which the river still perseveres. It is difficult here to decide how much confidence may be placed in the explanation, because of its giving reason for the location of certain streams, and how much doubt must be cast upon it, because it seems impossible and a sort of universal application.

39. *Migrations of the Atlantic-Ohio rivers.* There are ~~some~~ <sup>many</sup> shifted courses which cannot be definitely referred to any particular cycle and which may therefore be mentioned now. Among the greatest are those by which the divide between the Atlantic and the Ohio streams has been changed from its initial position on the great constructional Nittany highland and Bedford range. There was probably no significant change until after Newark deposition, for the branches of the Anthracite river could not have begun to push the divide westward till after the eastward flow of the river was determined; now, then, there does not seem to have been any marked advantage possessed by the eastward streams over the westward. But with the eastward escape of the Anthracite, it promptly found a shorter route to the sea and one that led it over alternately soft and hard rocks, instead of the longer course followed by the Ohio streams over continuous sandstones. The advantage given by the greater extent of soft rocks is indicated by the great breadth of the existing valleys in the central district compared with the less breadth of those in the plateau to the west. Consider the effect of this advantage at the time of the Jernsaw elevation. As the streams on the eastern slope of the Nittany divide had the shortest and steepest courses to the sea, they deepened their valleys faster than those on the west and acquired drainage area from them; hence we find reason for the drainage of the entire Nittany and Bedford district by the Atlantic streams as present. Various branches of what are now the Alleghany and Monongahela originally rose on the western slope of the dividing range. These probably reached the ~~Atlantic~~ <sup>ocean</sup> far to the east. ~~But Pennsylvania was too far to the west~~ <sup>for the rivers to reach the ocean</sup> ~~for the rivers to reach the ocean~~ <sup>before the great anticline was formed</sup> ~~but the summit antichlinal of Laurel ridge and Negro mountain farther west do not seem to have been strong enough to form a divide, for the rivers still traverse them.~~ Now as the headwaters

The main branch of the eastern slope of the Nittany-Bell range and passed the divide westward, they at last gained possession of the Siluro-Devonian monocline on its western side; but beyond this it has not been possible for them yet to go. As the stream cut down deeper and encountered the Medina and came near the core of the ridge, they made a passage through it; the Cambrian beds were discovered below and a valley was opened on them as the Medina cover wore away. The most important point about this is that we find in it an adequate explanation of the opposite location of water-gaps in pairs, such as characterize the branches of the Juniata below Tyrone and again below Bedford. This opposite location has been held to indicate a antecedent origin of the river that passes through the gaps, while gaps formed by self-developed streams are not thought to present such correspondence (Hilber). Yet the special case of paired gaps in the opposite walls of a breached anticline is manifestly a direct sequence of the development of the Juniata headwaters. The settling down of the main Juniata on Jack's mountain anticline below Hampton is another case of the same kind, in which the relatively low anticlinal crest is as yet not widely breached; the gaps below Bedford stand apart, as the crest is there higher, and hence wider opened; and the gaps below Tyrone are separated by some ten or twelve miles.

When the headwater streams captured the drainage of the Siluro-Devonian monocline on the western side of the ancient Nittany-Bell range, the main branch of the river was growing like a well trained grape vine. Most of this valley has been acquired by the west branch of the Susquehanna, probably because it traversed the Medina beds less often than the Juniata. For the same reason, it may be, the West Branch has captured a considerable area of plateau drainage that must have once belonged to the Ohio, while the Juniata has none of it, but if so, the capture must have been before the Tertiary cycle, for since that time the ability of the West Branch and of the Juniata as regards such capture appears about alike. On the other hand Castelman's river, a branch of the Monongahela, still retains the drainage of a part of the Siluro-Devonian monocline at the southern border of the State, where the Juniata headwaters had the least opportunity to capture it; but the change here is probably a recent one, not prevented, however, because the Juniata was so nearly yet to be driven back to the Allegheny from the frontal land of the plateau.



40. *Other examples of adjustments.*—Other examples of similar adjustments are found around the Wyoming basin, fig. 23.



FIG. 23

Originally all these streams ran centripetally down the enclosing slopes, and in such locations they must have cut gashes and breaches in the hard Carboniferous beds and opened low back country on the weaker Permian. Some of the existing streams still do so, and these are precisely the ones that are not easily reached by diverters. The Susquehanna in its course on side of the basin has sent out branches that have headed all the centripetal streams within reach; where the same river enters the basin, the centripetal streams have been shortened if not completely headed. A branch of the Delaware has captured the heads of some of the streams near the eastern end of the basin. Elsewhere, the centripetal streams still exist of good length. The contrast between the persistence of some of the centripetal streams here and their peripheral diversion around Broad Top is a consequence of the difference of altitude of the old lake bottoms in the two cases. It is not to be doubted that we shall become acquainted with many examples of this kind as our intimacy with rivers increases.

41. *Events of the Quaternary cycle.*—The brief quaternary cycle does not offer many examples of the kind that we have considered, and almost none found are of small dimensions. The only capturing stream that should be mentioned has lately been

\* Science, xii, 1899, 105.

ways have been affected by the same forces, and the relations of the greater streams in earlier cycles.

The character of the streams and their valleys as they now exist is strikingly dependent in many ways on the relation of the incipient primary cycle to the larger cycles of the past. No lakes occur, exception being made only of the relatively small ponds due to drift obstruction within the glaciated area. Waterfalls are found only at the headwaters of small streams in the glaciated area, every larger stream forming a continuous series of larger streams that have been thrown from their pre-glacial courses by drift barriers, and which are now in a very immature state on their new lines of flow. The small valleys of this cycle are shallow and narrow, always of a size strictly proportional to the volume of the stream and the hardness of the enclosing rocks, exception being made only in the case of post-glacial gorges where streams have been captured from older pre-glacial courses.

The terraces that are seen, especially on the streams that flow from the glaciated district, are merely a subordinate and subordinate complication of the general development of the valleys. In the region that has been here considered, the streams have been seldom much displaced from their pre-glacial channels; but in the northwestern part of the State, where the drift in the valleys seems to be heavier, more serious disturbances of pre-glacial courses is reported. The facts here referred to in regard to lakes, falls, gorges, terraces and displaced streams are to be found in the various volumes of the Second Geological Survey of the State,\* in regard to the terraces and the estuarine deflection of the Delaware and Susquehanna, reference should be made also to Meade's studies.†

4. *Doubtful cases*.—It is hardly necessary to state that there are many facts for which no satisfactory explanation is found under the theory of adjustments that we have been considering. Some will certainly include the location of the Susquehanna on the points of the Potomac arches under the category, a. I do not feel that such a location savors of an antecedent origin. It seems probable that the examples of the post-glacial waterfalls found on certain streams, for example, the four gaps cut in the

\* Especially Carr, Reports I., II., White, Reports G., H., Lewis, Report Z.

† Amer. Journ. Science, xxxv, 1888, 467, 448. Seventh Annual Rep. U. S. G. S., 1888, 645.

two pairs of Pocono and Portsville outcrops at the west end of the Wyoming syncline, and the three gaps where the Little Susquehanna crosses the Allegheny at Tyrone and Bedford have already been sufficiently explained. The location of the upper North Branch of the Susquehanna is also unrelated to processes of adjustment as far as I can see them, and the great area of plateau drainage that is now possessed by the West Branch is certainly difficult to understand as the result of conquest. The two independent gaps in Tussey's mountain, maintained by the Juniata and its Frankstown branch below Tyrone are curious, especially in view of the apparent veering of the branch to the main stream on the upper side of Warrior's ridge (Oriskany), just east of Tussey's mountain.

13. *Complicated history of our actual rivers.*—If this theory of the history of our rivers is correct, it follows that any one river as it now exists is of so complicated an origin that its development cannot become a matter of general study and must unhappily remain only a subject for special investigation for some time to come. It was my hope on beginning this essay to find some teaching sequence of facts that would serve to relieve the usual routine of statistical and descriptive geography, but this is not the result that has been attained. The history of the Susquehanna, the Juniata, or the Schuylkill, is too involved with complex changes, if not enshrouded in mystery, to become intelligible to any but advanced students, only the simplest cases of river development can be introduced into the narrow limits of ordinary instruction. The single course of an ancient stream is now broken into several independent parts, witness the disjointing and diversion of the original Juniata, which, as I have supposed, once extended from French Top lake to the Catawissa basin. Now the upper part of the stream, representing the early Broad River stage, is reduced to small volume in Aughwick creek; the continuation of the stream to Lewisstown is first set back some of its original head branches and is then diverted to another system; the beheaded portion, now represented by Middle creek is diverted from its course to the Catawissa basin by the Susquehanna, perhaps the Catawissa of the present day represents the reversed course of the lower Juniata where it joined the Attumate. This unservicably complicated statement is not much simplified if instead of beginning with an original stream and searching out its present disjointed parts, we trace the composition of a single

existing stream from its once independent parts. The drainage of to-day consists of headwaters acquired from Ohio streams; the lake in which the river once gathered its upper branches is now drained and the lake bottom has become a mountain top, the

a short course towards Louisiana nearly coincides with the original direction of the stream, not to confound this with a previous adjustment is to lose the true significance of river history. The course is the product of diversions at least at two spots and certainly in several places; and where the river now joins the Susquehanna it is suspected of having a superimposed course altogether different from the rest of the stream. This is too complicated, even if it should ever be demonstrated to be wholly true, to serve as material for ordinary study, but as long as it has a savor of truth, and as long as we are ignorant of the whole history of our rivers, through which alone their present features can be rightly understood, we must continue to search after the natural processes of their development as carefully and thoroughly as the geologist searches for the links missing from his scheme of classification.

44. *Pennsylvania Characteristics.*—It is in view of these doubts and uncertainties that I feel that the history of our rivers is not yet written. Yet for numerous occurrences of actual and deductive facts—proof so definite and in some cases so remarkable that they cannot be neglected, as they must be if we should adhere to the antiquated theories of the river courses.

The method adopted on an early page therefore seems to be justified. The provisional system of ancient drainage, arranged on fig. 2, does appear to be sufficiently related to the streams of to-day to warrant the belief that most of our rivers took their first courses between the primitive folds of our mountains, and that from that distant time to the present the changes they have suffered are due to their own interaction—in their own mutual adjustment more than to any other cause. The Susquehanna, Seneca II, Lehigh and Delaware are exceptional, composite and largely composite rivers, of repeated mutative adjustment. The middle Susquehanna and its branches and the upper portions of the Seneca II and Lehigh are descendants of original Pennsylvania rivers consequent on the construction topography of that time. Newark depression reversed the flow of some of the transverse streams, and the spontaneous outflow of adjustments from immen-

ture to mature courses in the several cycles of development are so numerous and extensive that, as Low truly says, the initial drainage has almost disappeared. The larger westward-flowing streams of the plateau are of earlier, Carboniferous birth, and have suffered little subsequent change beyond a loss of head waters. The lower courses of the Atlantic rivers are younger, having been much shifted from their Permian or pre-Permian courses by Newark and Cretaceous superimposition, as well as by recent downward deformation of the surface in their existing estuaries. No recognizable remnant of rivers antecedent to the Permian deformation are found in the eastern part of the State; and with the exception of parts of the upper Schuylkill and of the Susquehanna near Wakes-Barre, there are no large survivors of Permian consequent streams in the ordinary meaning of the term "consequent." The shifting of courses in the progress of mature adjustment has had more to do with determining the actual location of our rivers and streams than any other process.

Harvard College, June, 1888

## TOPOGRAPHIC MODELS

BY CONRAD MINDELEFF

OF the many methods by which it has been sought to represent the relief of a country or district, only two have been at all widely used. These methods are, in the order of their development, by hachured and by contoured maps. Both have advantages and both have serious disadvantages. Without entering into the controversy that is even yet raging over the relative merits of the two systems, some slight notice of what each claims to accomplish is necessary.

The representation of relief by hachures is a graphic system, and in the best examples we have is an attempt to show, upon a plane surface, the actual appearance of a given area under given conditions of lighting,—as in the Dufour map of the Alps. Of course certain details that would really disappear if the assumed conditions were actual ones, must be shown upon the map,—so that it is, after all, but a conventional, representation. The very best examples are, for this and other reasons, unsatisfactory, and far more so than the case to the vastly larger class of mediocrity and poor work.

The contour system represents relief by a series of lines, each of which is, at every point throughout its length, at a certain stated elevation above sea-level, or some other datum plane; in other words, each contour line represents what would be the water's edge, if the sea were to rise to that elevation. It preserves the advantage of great clearness, but falls to a large degree in the representation of surface detail; moreover, one must have considerable knowledge of topography, in order to read the map correctly.\*

To those who must give first place to the quantity of relief rather than the quality, as, for example, the geologist or the engineer, a contoured map is now considered essential. On the other hand, where quality of relief is the prime consideration, and the quantity a secondary one, as, for example, for the use of the army, a hachured map is considered the best. The method

\* For specimens of representation of the same subject in different series, in both the hachured and contour systems, see plates from "E. L. Offer's *Topographische Atlas*."

of hachures may be roughly characterized as a graphic system with a conventional element, and the contour method as a conventional system with a graphic element, -for if the contour interval is small enough a sort of shading is produced which helps considerably the idea of relief.

In addition to these two great systems, with which everyone is more or less familiar, there is another method of representing a

country which although it is a new view, has not received the attention it deserves.

It is the representation of a country by a model in relief. Certain striking advantages of models over maps of all kinds are, indeed, so apparent that one almost loses sight of such slight disadvantages as can, of course, be urged against them. In the graphic representation of the surface they are far superior to the hachured map, and they have the further advantage of expressing the relative relief, which the hachured map fails to do, except in a very general way. They have also the advantage of showing actual shadows, exactly as they would be seen in a bird's-eye view of the district, instead of more or less conventional ones, and are, consequently, more easily comprehended by the layman, without becoming any less valuable to the skilled topographer. In short, they combine all the graphic features of a hachured map with all the advantages of the best class of contoured maps, and in addition they show more of the surface detail, upon which so much of the character of the country depends and which is very inadequately expressed by hachures and almost completely ignored in a contoured map of large interval. The contours themselves can be made to appear upon the model, very easily and without interfering with other features.

The uses of models are many and various. Within the past few years their usefulness has been much extended, and now that they are becoming better known, will probably receive a still further extension. To the geologist they are often of great value in working out the structure of complicated districts, for the reason that so many important structural relations can be presented to the eye at a single glance. Similarly, for the graphic presentation of results there is no better method, as the geology, the surface geology, and any number of sections can be shown together and seen in their proper relationship. To the engineer an accurate model is often of the greatest assistance

and set out his problems, and it is simply invaluable to explain the details of a point to anyone who has little or no technical training; for, as has been stated, a model is easily comprehended by anyone, while more or less technical knowledge is required for the proper understanding of even the best maps.

I might go on cataloguing in detail the many uses to which models may be put, but shall now mention only one more—perhaps the most important of all—their use in the education of the young. No method has yet been devised that is capable of giving so clear and accurate a conception of the principles of physical geography as a series of well selected models, models have, indeed, already been used for this purpose, but unfortunately their great cost has prevented their general use in schools. Since, however, the study of geography has been placed upon a new basis and a new life has been infused into it, many men have given their attention to the subject of models and have experimented with a view to cheapen the cost of reproduction, which has hitherto prevented their wide distribution; and probably this object is well soon to be remedied. The ability to read a map correctly,—to obtain from a study of the map a clear conception of the country represented,—is more uncommon than is usually supposed. Some of the recent methods of teaching geography are intended to cultivate this very faculty, but it is doubtful whether there is any better method than that which consists in the study of a series of good models in conjunction with a series of maps, all on the same scale and of the same area. The value of a series of good models in teaching geology is so apparent that it need only be mentioned. It is often for reasons stated above, far more valuable even, than field instruction.

For the construction of a good relief map the first requisite is a good contoured map. To this should be added, when possible, a good bathymetrical map, upon which the elevations of the principal points are stated,—if the interval in the contoured map is a large one,—and as much material in the way of photographs and sketches as it is possible to procure. The modeler should, moreover, have some personal acquaintance with the region to be represented, or, failing that, a general knowledge of topographical forms, and at least a clear conception of the general character of the country which he seeks to represent. This is very important, for it is here that many modelers fail: the mechanical portion of



the work any ordinary intelligent person can do. A model may be as accurate as the map from which it is made, every contour may be placed exactly where it belongs, and yet the resulting model may be, indeed, often is—"flat," expressionless, and unsatisfactory. Every topographer in creating his map is compelled to generalize more or less, and it is fortunate for the map if this be done in the field instead of in the draughtsman's office. But

geographers differ among themselves: there may be, and often is, considerable difference in two maps of the same region made by different men; in other words, the "personal equation" is a larger element in a map than is usually supposed. This being the case, there is something more required in a modeler than the mere transferring of the matter in the map,—giving it three dimensions instead of two: he must supply through his special knowledge of the region (or, failing that, through his general knowledge) certain characteristics that do not appear upon the map, and it may, so far as it is necessary, certain generalizations of the topographer and draughtsman. This artistic or technical skill required correctly to represent the *individuality* of a given district is especially important in the modeler; it is more important, perhaps, in small scale maps of large districts than in large scale maps of small ones,—for in the latter the generalizing process has not been carried so far, and the modeler in *even* of the *small* scale often preserves much of the detail.

The methods by which relief maps are made have always received more attention than would, at first sight, appear to be their proper proportion. It may be due, however, to the difficulty of applying any test to determine the accuracy of the finished model, and perhaps also to the general impression that any one can make a relief map, and so he can, though of course there will be a wide difference in the value of the results. Some, indeed, have devoted their attention to methods exclusively letting the result take care of itself,—and the models show it. There is no more reason why a modeler should be looked down upon as a dilettante, than that a chemist, artist, or a rhetorist, or anyone engaged in technical work, should be so; though in some cases he might be required, as the chemist is, to show his model as well as his results.

One of the earliest methods, with any pretension to what we may term mechanical perfection, is that described by the Messrs. Alabach in a paper on "The construction of maps in relief," read

before the American Institute of Mining Engineers in 1887. The method was published in 1888. Upon a contoured map as a basis cross-section lines are drawn at small and regular intervals, and, if the topography be intricate, corresponding lines at right angles. The sections are carved out of some solid material of some suitable material, such as cardboard or metal, and cut down to the surface line,—the strips themselves thus forming the cross-sections. These cross-sections are mounted upon a suitable base-board, and the cavity or boxes are then filled up with some easily carved material, such as plaster or wax. The top is then carved down to the form of the country or district,—the necessary guidance being obtained by the upper edges of the strips that form the cross-sections. It will be readily seen that this method is a very crude and laborious one. It necessitates in the first place a good contoured map upon which to draw the sections, but sacrifices much of the advantage thus gained because only a number of points on each section line are used, instead of the entire line. It is no better, although actually more laborious, than the later method of driving contour pins (whose height above a base-board may be accurately measured,) along the contour lines, and then filling in. A slight modification of the latter method can be used to advantage when no contoured map is available, and when the points whose elevation is known are not numerous enough to permit the construction of one. In this case the only control that can be secured is by means of a number of pins driven into the base-board at those points whose elevation is known. The remainder of the map is then sketched in. This method is perhaps as satisfactory as any, when the material upon the map is ready. Another method, however, growing out of the same want, use of material, is in some cases to be preferred, especially for large models. The map is enlarged to the required size, and a tracing of it is mounted upon a frame. Another deep frame, just large enough to contain the material, tracing, is made, and laid upon a suitable base-board upon which a copy of the map has been mounted. Upon this base-board the model is then commenced, in clay or wax. The low areas are modeled first,—horizontal control being obtained by picking through the mounted tracing of the map with a needle point and vertical control by measuring down from a straight edge sliding on the top of the deep frame. This system is rather crude, and only useful where the material upon the map is very expensive, but it gives excellent control.

A method used by Mr. F. H. King in the preparation of his large map of the United States is described by him in a letter to Messrs. Harden, and published by them in the place mentioned. A solid block of plaster is used; the contour map being transferred to it—and the plaster is carved down to produce a series of steps like those made by building up the contour. The shoulders are then carved down to produce a continuous surface. This method is one of the best of those that require carving instead of modeling.

Many other methods of producing relief maps might be mentioned, but, as most of them have been used only to make special models, they need not be described. The method that has been more used than any other at H remains to be described. It is this which the writer has used almost exclusively, and consists in building up the model and modeling the details, instead of carving it. It is a maxim of the modeler that the subject should be built up as far as possible, should be produced by adding bits of clay or wax, or other material, and not by carving away what is already on—by addition and not by subtraction. This may be illustrated by a reference to the methods of the sculptor. The bust or figure, or whatever the subject may be, is first modeled in clay or wax, from this is made a plaster mould, and from this mould a plaster cast is taken. The cast is called the original, and the finished production, whether in marble, bronze, or any other hard substance, is simply a copy of this original. No one ever attempts to produce the busted bust or figure directly from the object itself. Even where the artist has for a guide a death mask, the procedure does not change. The bust is first made in clay and this clay model as a rule, contains all the detail which subsequently appears in the finished bust. It seems strange, therefore, that the relief map maker should use a method which the sculptor, with infinitely more skill and judgment, is afraid to use; and that on subjects that do not differ as much as might be imagined.

The contour interval to be used depends on the use to which the model is to be put. It is not always necessary to carry into the model all the contour lines upon the map. I may go farther and say that it is not always desirable to do so. The number to be used depends to some extent on the skill of the modeler. As already stated, the contours are only a means of control, and one modeler requires more than another. To build into a model every

contour in a contoured map of ten foot interval is a very laborious proceeding, and not worth the time it takes, as in nine out of ten maps of such interval only the fifty foot or the one hundred-foot curves are definitely drawn, the intermediate lines being merely filled in. This thing can be done as well, or better, by the ruler.

The question as to the proper amount of exaggeration to be given the vertical scale, as compared with the horizontal, is the question about which has raged most of the controversy connected with relief map making. This controversy has been rather bitter; some of the exponents of vertical exaggeration going to the length of saying that no exaggeration is necessary, and that "no that would distort or exaggerate the scale of any thing whatsoever." On the other hand the great majority of those who have made relief maps insist upon the necessity of more or less exaggeration of the vertical scale—generally more than seems to be necessary, however.

An increase of angle of slope accompanies an vertical exaggeration, and this is apparent even in models in which the vertical scale is not at all vertically exaggerated. It produces a false effect by diminishing the proportionate width of the valleys, and by making the country seem much more rugged and more mountainous than it really is. A secondary effect is to make the region represented look very small—a false idea of the extent of the country being lost. This can be illustrated better than described. The King model of the United States is an example of one extreme; it is worthy of note that no examples of the other extreme—too little exaggeration—are known.

In small-scale models of large districts some exaggeration of the vertical scale is necessary in order to make the relief apparent, but the amount of this exaggeration is often increased much beyond what is essential. The proportion of scales must depend to a large extent on the character of the country represented, and on the purposes for which the model is made. It has been suggested by a writer, quoted by the Messrs. Harden, that the following exaggeration would afford a pleasing relief: "For a map, scale 6 inches to 1 mile: if mountainous, 1.5, if only hilly 1.2; if gently undulating, .5-1. For smaller scales, except for sea-level purposes, the vertical scale should be increased in proportion." For a tract consisting wholly of mountains no exaggeration is necessary." I know of no country of such a charac-



as his idea of it would be if he were on the real ground itself (are should be taken, however, not to make the scales so disproportionate as to be violative to mental impressions. " . . . indeed, prominent or important features when they were here it may be still more effectively shown by systematic exaggeration in the vertical scale." The fallacy of this argument is obvious. It assumes that the object of a model is to show the country as it appears to one passing through it, and not as it really is—and there is often a very wide difference between the two. The impression derived from passing through a country is, if I may use the term, a very large-scale impression, as any one who has tried it can certify. It is certainly a mistake to attempt to reproduce this impression in a small-scale model, with the aid of vertical exaggeration. Even if the principle were a good one, its application would be very limited. It could only be used in large-scale models; to apply it to a model of a large area—the United States, for example—would be utterly absurd.

The method referred to as being now generally in use may be briefly described as follows: requiring, a good contour map; a hachured map in addition, if possible; a clear conception on the part of the modeler of the country to be represented; and a fair amount of skill. Materials, a base-board of wood or other suitable material; card-board or wood of the thickness required by the contour interval and the scale; modeling wax or clay. Procedure, reproduce the contours in the wood or other material; mount these upon the base-board in their proper relative relationship; then fill in the intervening spaces, and the space above the topmost contour, with the modeling material.

In a series of models of the Grand Divisions of the earth, made about a year and a half ago, the contours of card-board were made as follows: the map was photographed up to the required scale, and as many points were made as there were contour intervals to be represented—in a model of the United States of 1,000 feet contour interval there were fourteen points. Thirteen of these were mounted upon card-board of the exact thickness required by the vertical scale, and one upon the base-board. All large paper companies use a micrometer gauge, and card-board can easily be obtained of the exact thickness required—even to less than the thousandths part of an inch. The lowest contour was then sawed out upon a steel saw, and placed upon the card-board just as the others were placed, and the model was

It was repeated with a similar result, and the result was not better. At this stage the model is prepared by a series of steps, each step representing a rise corresponding to the contour interval. The advantage of this method is, the fact that even the greatest distortion in making the plan of the model will be a negligible distortion, owing to the stretching of the paper in the first step is not large, and the distortion in fitting the contours, if care be exercised in having the grain of the paper in the same direction in all the pieces, the distortion in fitting the contours will be much reduced, but the distortion in one direction will remain. In our experience the distortion amounts to about two per cent; in other words, a model that should be fifty feet high will in reality be fifty-one feet high, but as this error is distributed over the whole fifty feet it is not so great for an ordinary model. If greater accuracy be required, it can be secured by transferring the contours to the cardboard by means of tracing paper. The great advantage of the plan of the model is the fact that when the model has been made, with the contours in position, it presents a copy of the relief, with all the details of drainage, etc., in position, instead of blank intervals between the contours. Such details and drainage are a great help in the subsequent modeling.

The next step in the process is to fill in with clay or wax the intervals between the contours. I have always found wax more convenient than clay for this purpose as, unless the surface coating is a thick one, the clay is difficult to keep moist. To obviate this difficulty we have used clay mixed with glycerine instead of water; this, of course, does not become dry, but the material is, at its best, unsatisfactory. The filling-in process is the most important one in relief map making, for it is here that the modeler must show his knowledge of, and feeling for, topographic forms. Some models seem to have been constructed with the idea that when the contours have been accurately placed the work of the modeler is practically done. This is a great mistake. The cardboard contours are only a means of control, occupying somewhat the same relation to the relief map that a core or base of bricks, or a frame of wood does to other construction, for example, an architectural ornament or a bust. It is sometimes necessary to cut away the contour card; for, as the contours are placed, a line is not necessarily perfect, as I

a comb or a frequently carried ruler as a razor—instead of following it up, as it would be if the rills were on a larger scale. Such generalizations of a topographer's proper is a map. At the same scale, we expect more detail in a model. The modeler must have judgment enough to know enough to read between the lines, and to undo the generalizing of the topographer and enough sense to say "no" to the map, and to say "yes" from the map. This can be done without materially affecting the accuracy of the model, considered even as a copy of the contoured map.

The contours of card board or other material are, let me repeat, only a means of *aid*. The person who makes a model by the way, yet to be evolved—would be able to make an accurate model that would stand on the same way that the topographer's are made. So, for example, a flower on an acre, an architectural ornament, or any other subject in low relief, where the object sought is artistic effect, and great accuracy is not a desideratum. It is the converse of this idea that has produced the numerous models that one sees; accurate enough, perhaps, but wholly expressionless and absolutely without feeling. This is the great fault of these models, and it is to be remedied. The models in wood and plaster are a whole lot better. It is then necessary to say "no" to them, and what little character they might otherwise have had is completely obliterated by the sand-paper. Such models almost invariably *look* wooden. Let the modeler, then, have a clear conception of his subject and not depend wholly on the contours, and let him work out this conception in his model, "controlled" and helped by the contours, but not bound by them; the resulting model will thus be far more satisfactory and a far better representation of his subject, in other words, it will be more life-like—in *expression*.

The model, provided it be not of clay, is sometimes used in the state in which it is left when finished. It is much more common, however, to make a plaster mould, and from this a plaster cast. For this purpose a moulder is usually called in; but moulders as a rule are ignorant men, accustomed to one line of work only, and the result is not always satisfactory. It is much better for the modeler himself to do this work, though to obtain good results from plaster it is necessary to know the material thoroughly, and this knowledge comes only from experience. The mould is generally made quite heavy, in order to stand the subsequent hard treatment that it may receive, and should be retouched and thor-



oughly dried before being prepared for the cast. The method used by some modelers of placing a frame about the model and pouring in the plaster, filling the frame to the top, is a crude and very wasteful one and not at all to be recommended. In a mould of large size—say seven or eight feet square—it would require a derrick to move the mould. It is wholly unnecessary, as, with a small amount of care, a good mould can be made not more than an inch thick, or, at most, an inch and a half. The drying of the plaster is not a matter of great importance, but it is always desirable.

**Notes.**—An English method is to grease the form. French and Italian ones varnish the mould, and thus lose some of the finest detail and sharpness. This is unnecessary. The mould can be easily prepared with a solution of soap so as to leave nothing on the surface but a very thin coating of oil, which is taken up and replaced by the plaster of the cast. Of course, if the model has been sand-papered, no fine work in moulding or casting is necessary, as there is nothing to save. If the subject is a very intricate one, with "undercuts" (as they are called), it is customary to make a waste mould; as this is very seldom necessary in relief map work, however, the process need not be described.

To make the cast it is only necessary to repeat the processes used in making the mould. With great care and some skill a cast can be made of the same size or of a different shape as the original model. It is customary to make the cast very thick, and, consequently, very heavy; this is unnecessary. In our work we seldom make a cast thicker than one inch, and yet are never troubled with changes in the model after it is finished. Even in a very large cast (now in the National Museum, weighing nearly 1,500 pounds and presenting a surface of over 100 square feet, the average thickness is less than one inch, although it required over five barrels of plaster to make it. The cast is not dried, but is allowed to be treated as the original model, imperfections being carefully repaired. The surface, however, should be touched as little as possible, as the slight roughness of surface that comes from the original model, through the mould, is removed by any tooling. The roughness adds much to the effect of the model—a fact which has been often observed. It is sometimes desirable to emphasize it.

The proper way to paint a model is a matter that must rest principally upon the judgment of the modeler, depending on some

extent, and, on the use to which the model is to be put. The plan used is a *matte* finish, meaning a surface that does not reflect light. This has the advantage of preserving all the detail that comes from the model, but it has also the disadvantage of a surface easily soiled and impossible to clean. If the model is to be painted, the surface should be *semi-gloss*—in our practice we use a small amount of yellow with the white. This yellow is hardly appreciable by the eye, but its effect upon the photographic negative is quite marked. Yellow becomes green in a photograph, and a photograph of a model covered as described, a grey tint is given to the whole surface. The colors, however, are not pure white and black as in a photograph; but a light and a dark. There is another kind of photographic surface, photographing models, the surface should have a *semi-gloss* that would have no gloss, but at times, should have only what is known among painters as an *egg-shell gloss*. It is almost impossible at this day to photograph a model that has a *semi-gloss* surface. Any object for a model that is desired separate from the rest should be painted in a different color—the water, for example, should be painted a light blue, not a blue composed of indigo, however, or any of the gray blues, as these produce in the photograph a small grey, and are not pleasant to the eye. The most satisfactory color that we have used is a mixture of white, the color of the paper, with a very little—quite green—and white. This gives a color that is pleasant to the eye, and is very pleasing in a photograph and is well.

Models intended for exhibition are so to be painted—painted realistically. There is room here for an immense improvement in the usual practice, which is to paint the model either in some conventional color of light and dark, or to put a single flat tint upon it. If the model is to be colored conventionally it is, in my opinion, much better to use a flat tint, light in tone, and with a dead surface. The use of a variety of colors upon the face of a model is not a matter of course, especially if the model is to be used for a long time. For a reason, much better to have a single color, as a model can be used as a single duplicate representing topography only, colored realistically, if possible, and without lettering. Well-defined lines other than those pertaining to the model itself, such as for example, as those used to define the boundaries of geological formations, should not

be allowed upon a model when it is desired to bring out all the relief. The lettering on such models should be kept down as small as possible, or when it is desired to be small the letter is much the better method.

The cheap reproduction of models is the most important problem connected with the art of making models. If models are to be reproduced cheaply, they will never have any wide distribution and there will be but less incentive to the modeler. Various materials have been suggested and experimented on, but the models of the mountains that are made today are made of plaster of Paris. Although this material was the first to be used for this purpose, it has not yet been superseded. A plaster cast is easy, expensive as it is very good. The plaster gives an accurate copy of the original surface, particularly if the form given to the model is not too large. The weight is not too heavy, but it is not very strong. By the incorporation of the plaster of Paris with, or by lugging it with, of various kinds, it can be made to make very good casts at the same time, and, as the expense is increased rather than decreased by this method. Models made in this way, however, have the advantage that when broken the plaster does not fall out, it never falls out, and it is not so brittle as the other kind. The large cast in the National Museum, before referred to, was made in this way. It weighed nearly 2,000 pounds when boxed—not an easy thing to handle—but it was not injured. It was the same as a block without suffering any damage at all. This would hardly have been possible if the cast had been made from plaster alone.

Paper seems, at first sight, to be the most ideal material for the reproduction of models, but no one has succeeded well enough with it to bring it into use. Like nearly all those who have given this subject attention, I have experimented with paper, but the only positive result has been a loss of a large part of the confidence that I had had in the subject. If the material paper had been used extensively for large scale models, it would have been found that I have never obtained a satisfactory result with subjects on a low relief and fine detail. A paper cast may look well when first made, but it absorbs moisture from the atmosphere, and contracts and expands with the weather. The contraction is apt to flatten out the model, and the expansion may make it look cup.

Casts of models have been made in iron and brass, when an entire

perhaps for models of mountains and subjects of like character would hardly be applicable to small-scale models with fine detail; such casts require too much surface finishing. The material known as *lanterna*, Walton seems to me to be the ideal material for this purpose. It is tougher than rubber, will take the finest detail, and its surface can be treated in any way desired. Unfortunately the manufacture of models in this material would require expensive machinery, and is outside the scope of a modeling room. Should it ever become commercially advantageous, however, casts of a model of ordinary size, in every way equal to the original, can be turned out in this material at a very small cost.

It remains to speak of the reproduction of models by process-  
engravings—a method that will probably receive much more  
attention in the future than it has in the past. It is perhaps  
worth noting that the method of reproducing figures in wax  
and other subjects of the kind that I have just treated here  
has not been perfected until some future occasion.

Scale 1 inch = 4 miles



Hachure

"

Contour  
400 ft

Interval  
300 ft

Scale 1 inch = 1 mile



Hachure

Contour interval 40 ft

Contour interval 80 ft

Contour interval 20 ft

# HACHURED AND CONTOURED MAPS

REPRESENTATION OF A HILL ACCORDING TO THE TWO SYSTEMS  
AND ON DIFFERENT SCALES

From Supplement to Enithoffers Topographical Atlas  
by permission of Mr Enithoffers



# NATIONAL GEOGRAPHIC SOCIETY.

## ABSTRACT OF MINUTES.

### *October 5, 1888, Ninth Meeting.*

A paper was read entitled, "Topographic Models," by Mr. Clarence M. Felt. It appeared in the "National Geographic Magazine," Vol. I, No. 3.

### *October 19, 1888, Tenth Meeting.*

The attendance being very small, no paper was read.

### *November 2, 1888, Eleventh Meeting.*

The paper of the evening was entitled, "Surveys, their Kinds and Purposes," by Mr. Marcus Baker. The paper was discussed by Messrs. Ogden, Gould, Low, Gannett and Baker. Published in "Science," Vol. XII, No. 304.

### *November 16, 1888, Twelfth Meeting.*

A paper was read by Mr. L. Gannett, giving certain "Facts and Statistics of the Marine Survey of the United States," the map of that date recently prepared by the United States Geological Survey. A discussion followed which was participated in by Messrs. Baker, Johnston, Bernow, Weed, and the author. A second paper entitled, "Something about Trinidad," was read by Lieut. J. P. Hawley, U. S. Signal Corps.

### *November 30, 1888, Thirteenth Meeting.*

The annual reports of our Presidents Herbert G. Ogden and Gen. A. W. Greeley were delivered. Published in the "National Geographic Magazine," Vol. I, No. 4.

### *December 20, 1888, Fourteenth Meeting.*

Held in the Law Lecture Room of the Columbian University. The President delivered his Annual Address, entitled, "Africa." Published in the "National Geographic Magazine," Vol. I, No. 5.

*December 24, 1889, Fifteenth Meeting.*

The Society met in the Society Hall of the Cosmos Club, President Hubbard in the chair. Owing to the absence from the city of the Secretaries, Mr. O. H. Tittmann was requested to act as Secretary of the meeting. The minutes of the first and second meetings were read and approved. The report of the Secretaries was read, in their absence, by the temporary Secretary, and was approved. The Treasurer's report, showing a balance of \$1,000.00, was read and approved. The report of the auditing committee was read and approved.

The Executive Committee met in the evening, and a resolution of two of the managers, Messrs. W. D. Johnson and Henry Mitchell, had been filed by the Board on the 15th of November, by the election of Messrs. O. H. Tittmann and C. A. Kenniston, and that a vacancy caused by the resignation of Vice-President John R. Bartlett, had been filled by the election of Lieut. George L. Dyer, on November 30th.

The Society then proceeded to the election of officers, with following result:

*President*—HARRISON C. HUBBARD.

*Vice-presidents*—HERBERT G. COHEN, 1st; GEORGE L. DYER, (and A. W. ORRILL, [ret.] C. HART MERRICK, [life]; A. B. THOMPSON, [ret.]

*Treasurer*—CHARLES J. HILL.

*Recording Secretary*—HENRY CLANNETT.

*Corresponding Secretary*—GEORGE KENNAN.

*Managers*—CLEVELAND ABBE, MARION HARRIS, ROGERS BIRNEY, JR., R. BROWN GIBBS, W. B. POWELL, J. C. WELLS, C. A. KENNISTON, O. H. TITTMANN.


*January 11, 1890, Sixteenth Meeting.*

The paper of the evening was entitled "The Great Plains of Canada," and was presented by Professor C. A. Kenniston, of Howard University.

*January 25, 1890, Seventeenth Meeting.*

The paper of the evening was entitled, "Irrigation in California," by Mr. William Hammond Hall, State Engineer of California. To be published in the "National Geographic Magazine," Vol. I, No. 4.





11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 10



*February 8, 1880, Eighteenth Meeting.*

The following papers were read by Prof. W. M. Davis, of Harvard University: "Topographic Models," and "Certain Features of the Rivers of Pennsylvania." Published in the "National Geographic Magazine," Vol. I, No. 3.

*February 22, 1880, Nineteenth Meeting.*

The paper "The River of the South," "Flooding about Asheville, N. C.," by Mr. Bailey Willis. The paper was illustrated by various sketches and lantern slides. Discussion followed, which was participated in by Messrs. Baker, Merriam and Moore. To be published in the "National Geographic Magazine," Vol. I No. 4.

*March 8, 1880, Twentieth Meeting.*

The following amendments to the By-Laws were adopted  
[For Article VI substitute the following]:

ARTICLE VI

MEETINGS.

"Regular meetings of the Society shall be held on alternate Fridays, from November until May, and excepting the annual meeting, they shall be devoted to communications. The Board of Managers shall, however, have power to postpone or omit meetings, when deemed desirable. Special meetings may be called by the President.

"The annual meeting for the election of officers shall be the next regular meeting in December.

"The meeting preceding the annual meeting shall be devoted to the President's annual address.

"The reports of the retiring Vice-Presidents shall be presented at the meetings in January.

"A quorum for the transaction of business shall consist of twenty-five active men here."

In Article V, the words "and the first meeting shall be held immediately after the first paragraph of the article"

"The dues of members elected in November and December shall be credited to the succeeding year."

The following papers were presented: "A Trip to Panama and Darien," by Mr. R. L. George, and "Survey of Mason and Dixon's Line," by Mr. Mark B. Kerr.

A Trip to Panama and Darien, to be published in the "National Geographic Magazine," Vol. I, No. 4.

*March 22, 1889, Twenty-first Meeting.*

The paper of the evening was entitled, "Recent Events in the U. S. of Columbia," by Mr. W. E. Curtis. The discussion which followed was participated in by Messrs. Baker, Gamett, and others.

*April 5, 1889, Twenty-second Meeting.*

The paper of the evening was entitled, "House Life in Mexico," by Mr. A. B. Johnson.

*April 19, 1889, Twenty-third Meeting.*

This meeting was devoted to papers upon the Samoan Islands. The following programme was presented:

"Samoa; the General Geography and Hydrography of the Islands and Adjacent Seas," by Mr. Everett Hayden.

"Climate," by Prof. Cleveland Abbe.

"Narrative of a Cruise Among the Islands," by Capt. R. W. Meade, U. S. N.

"The Home Life of the Samoans and the History of the Islands," by Mr. W. E. Safford, U. S. N.

*May 3, 1889, Twenty-fourth Meeting.*

The paper of the evening was entitled, "Across Nicaragua with Trent and Machete," by Mr. R. E. Peary, U. S. N. To be published in the "National Geographic Magazine," Vol. I, No. 4.

*May 17, 1889, Twenty-fifth Meeting.*

The paper of the evening was entitled, "The Krakatau Eruption," by Dr. A. Graham Bell. The paper was discussed by Captain C. E. Dutton.

*(Translated by Mr. R. G. Lorch)*

## INTERNATIONAL LITERARY CONTEST

To be held at Madrid, Spain, under the auspices of the Commission in charge of the celebration of the Fourth Centennial Anniversary of the Discovery of America.

### PURPOSE

The work for which a prize is offered is to be a prose essay, a true historic picture giving a just estimate of the grandeur of the occasion to be celebrated.

So much has been written on this subject since the opening of the XVIth century that it would seem difficult to say anything new and good. Perhaps the details, perhaps the circumstances in the life and work of Columbus are worthy of no little research, but already the Royal Academy of History is engaged in the task of collecting and editing the unedited or little known papers bearing on this question.

The book required by this contest must be of a different nature: it must be comprehensive and synoptic, and must be sufficiently concise without being either obscure or dry.

Although there is an abundance of histories of America, of voyages and discoveries, of geographical science, and of the establishment of Europeans in remote regions of the earth, there is no book that sets forth as it can be done the combined efforts of the nations of the Iberian peninsula, who, since the commencement of the XVth century, have, with a fixity of purpose and marvellous energy, in almost a single century of silent efforts brought about the exploration of vast continents and islands, traversed seas never before cut by Christian prow, and in emulous strife obtained almost a complete knowledge of the planet on which we live.

There is a growing interest in our century in these important events; not to mention the circumstantial evidence borne by the charts of 1475 and the semi-fabulous voyages, such as that of Donay Vasco and others less apocryphal though seldom mentioned, such as that of Ferrer, begun in 1434, when Dr. Esquivel reached Cay Baguer, discovered Guisea, and

1492 with Liengne's arrival at Barcelona after circumnavigating the globe.

In all this activity very little occurs by chance. The progressive series of geographic discoveries, due to persistent premeditation and not to accident, was inaugurated at Sagres by the illustrious Prince of Portugal, the Infante Dom João de Mascarenhas.

Well might Pedro Nunes claim that from that time forth the progress of cosmography was more rapid than that of any other science, "unless our mariners sailed away better instructed and provided with more accurate maps of Asia, Africa, and Europe than the things with which cosmographers supplied them."

The great aim of the Infante Dom João de Mascarenhas on the 12th of October, 1492, when Columbus was the first European to set foot upon the untrodden shores of the New World, was to inaugurate a new era in the history of geography, as part of a summation of efforts, a grand development of an idea, a purpose to explore and know the whole globe, to spread the name and the law of Christ together with the civilization of Europe, and to reap a harvest of gold, spices, and all the riches of the Indies. The Infante Dom João de Mascarenhas was not alone in this. He was joined by many other nobles, who in turn got them from Mussulmans.

Doubtless the moving cause, whose gorgeous banner so many men of our peninsula followed, was clothed in great sentiments, good or bad; their hearts were filled with religious fervor, thirst for glory, ambition, Christian love, cupidity, curiosity and violent dissatisfaction (even during the Renaissance), to seek and undergo real adventures that should surpass the vain, fruitless, and fanciful adventures of chivalry; and to make voyages and discoveries that should surpass all that had been known, many of which, recorded in classic histories and fables, were now disinterred by the learned.

What must be described is the complete picture in all its details, the whole scene, the magnificent meeting may be said to have been a grand assembly of the world, looking into the stories, voyages and happy adventures of Bartolomé Diaz, Gama,

Admiral Albuquerque, Cabral, Balboa, Magellan, Cortes, Pizarro, Orellana, and a host of others, do not dim the glory of the hero whose contemporary it is to be celebrated, even though it heighten and add greater lustre to the work of civilization begun by Portugal.

The book once vaguely outlined must also contain a compendious introduction, notices of voyages, ideas, and geographic progress up to the date of D. Henrique's establishment at Sagres, and an epilogue or conclusion of greater extent, in which are examined and weighed the changes and progress that our subject has made, collectively, in the civilization of the world—in the commerce, economics and politics of the peoples, in regard to the broad field opened to the intelligent activity of Europe, over which it could spread and dominate; the abundance of data

for the extension of our knowledge of Nature, the unravelling of her laws, and penetration of her mysteries.

The vast, elevated argument of the book requires it be a finished work of art—clear, concise and coherent of plan, by its plan and order, in sobriety and unity of style, whose nobility and beauty must be in simplicity of phrase, correctness of judgment and richness of thought.

It may be written in any of our great languages, work written in French, Spanish, Portuguese, English, Italian, French or Italian.

The tribunal that is to award the prize will be composed of two members of the R. Acad. of History, and one member from each of the Academies of Moral Sciences and Physical Sciences and Natural Sciences, and one member from the Academies themselves.

Furthermore, there will be included in the tribunal the diplomatic representative of every power whose subject or subjects wish to enter the contest, which is to be done through said representative or some person duly appointed to act in his place.

The tribunal will elect its presiding officer and will decide on the merits of every work submitted, and the prize will be awarded as part of the vote.

Each work submitted in this contest must be neatly copied, in legible writing, on good paper, without the author's name but with a quotation to identify him afterwards.

Each author will, inclose a separate folded sheet on whose exterior is written his name, which he will sign and seal, and

sentence of his work, within, he will write his name and residence.

The folded sheets corresponding to the works that did not get a prize will be burnt publicly without being opened.

Though it is difficult to set a limit as to size, the works should not have more reading matter than is contained in two volumes of the shape and size of the complete works of Cervantes issued by Rivadeneyra in 1862-4.

If the plan or purpose of any of the works requires it, there may be added another volume of documents, maps, or other illustrations.

As it will take time to examine and judge the works, they should be sent to the Secretary of the R. Acad. of Hist. prior to January 1, 1892.

There will be first prize of 30,000 pesetas (\$5,700) and a second of 15,000 pesetas (\$2,850).

Besides this, each of the two successful authors will receive 500 copies of the printed edition of his work.

It rests with the Central Commission to determine the number of copies in the edition of each of the two prize works, and what disposition is to be made of the copies that are not given to the authors.

These (the authors) keep the right to re-print and to sell their works, and to translate them into other tongues.

The Commission, however, will have the right, if either or both prize works are in a foreign tongue, to have them translated and published in Castilian.

The Commission affix their seal to the preceding directions for the information of the public and government of those persons who desire to participate in the contest.

Madrid, June 10, 1890.

The Vice President, D. RAY OF VERA;  
Secretaries, JEAN VALERA, JEAN F. RIASO.









